



AAPG | BULLETIN

VOLUME 97 • NUMBER 11 • NOVEMBER 2013

REGULAR FEATURES

Previews.....	ii
Staff Editors.....	iii
Association	iv
House of Delegates.....	vi
Instructions to Authors	1848
Electronic Submission	1920
Geoscience Meetings Calendar	2058
Education Calendar	2084
Erratum	2102
Memorial	2126
Affiliated Societies	(viii)

ON COVER – Microbial bioherm (inset photo) and biostrome (background photo) in the Upper Cambrian Wilberns Formation along the Llano River near Mason, Texas. Bioherm was described by Wayne Ahr (1971) and is approximately 15 m thick. Photos by W.A. Morgan.

ACKNOWLEDGMENTS—

AAPG thanks
BP Brasil Exploration & Appraisal;
BP Upstream Technology;
Chevron;
Colorado School of Mines;
ConocoPhillips;
ExxonMobil;
Texas A&M University;
University of Kansas
and the AAPG Foundation
for financial support of this issue of the *Bulletin*.

Introduction: AAPG Hedberg Research Conference on Microbial Carbonate Reservoir Characterization—Conference summary and selected papers

Ernest A. Mancini, William A. Morgan, Paul M. (Mitch) Harris, and William C. Parcell 1835

RESERVOIR CHARACTERIZATION, SUBSURFACE STUDIES, AND OUTCROP ANALOGS

Sedimentary features, occurrence, and cyclicity of microbialite facies in the Roubidoux and Jefferson City Formations of Missouri and Kansas	1849
Chamandika Warusavitharana and William Parcell	1849
Sponge-microbial mound facies in Mississippian Tuscarria Limestone, Walker County, Alabama	
David C. Kopaska-Merkel, Steven D. Mann, and Jack C. Pashin	1871
Calcisponge-microbialite reef facies, middle Permian (lower Guadalupian), northwest shelf margin of Permian Basin, New Mexico	
Gregory P. Wahlman, David M. Orchard, and Govert J. Buijs	1895
Upper Permian (Zechstein) microbialites: Supratidal through deep subtidal deposition, source rock, and reservoir potential	
Miroslaw Slowakiewicz, Maurice E. Tucker, Richard D. Pancost, Edoardo Perri, and Michael Mawson	1921
Lithofacies, stable isotopic composition, and stratigraphic evolution of microbial and associated carbonates, Green River Formation (Eocene), Piceance Basin, Colorado	
J. Frederick Sarg, Suriajin, Kati Tänavsuo-Milkeviciene, and John D. Humphrey	1937
Vertical and lateral distribution of lacustrine carbonate lithofacies at the parasequence scale in the Miocene Hot Spring limestone, Idaho: An analog addressing reservoir presence and quality	
Kevin M. Bohacs, Kathryn Lamb-Woziak, Timothy M. Demko, Jason Eleson, Orla McLaughlin, Catherine Lash, David M. Cleveland, and Stephen Kaczmarek	1967
Topographic and sea level controls on oolite-microbialite-coralgal reef sequences: The terminal carbonate complex of southeast Spain	
Robert H. Goldstein, Evan K. Franseen, and Christopher J. Lipinski	1997

RESERVOIR MODELING

Reservoir analog model for oolite-microbialite sequences, Miocene terminal carbonate complex, Spain	
Christopher J. Lipinski, Evan K. Franseen, and Robert H. Goldstein	2035
Reservoir characterization, modeling, and evaluation of Upper Jurassic Smackover microbial carbonate and associated facies in Little Cedar Creek field, southwest Alabama, eastern Gulf coastal plain of the United States	
Sharbel Al Haddad and Ernest A. Mancini	2059

PETROPHYSICS AND GEOCHEMISTRY

Three-dimensional pore connectivity evaluation in a Holocene and Jurassic microbialite buildup	
Marcelo F. Rezende, Sandra N. Tonietto, and Michael C. Pope	2085
Porosity in bacterially induced carbonates: Focus on micropores	
Henry S. Chafetz	2103
Ordered low-temperature dolomite mediated by carboxyl-group density of microbial cell walls	
Paul A. Kenward, David A. Fowle, Robert H. Goldstein, Masato Ueshima, Luis A. González, and Jennifer A. Roberts	2113