

ПИ
A60/6

ANNALS OF BOTANY


Founded 1887

EDITORIAL OFFICES IN UK, AUSTRALIA, CANADA, CHINA, JAPAN AND MAINLAND EUROPE

Volume 111

Number 5

May 2013



www.aob.oxfordjournals.org

ISSN 0305-7364 (PRINT)

ISSN 1095-8290 (ONLINE)

ANNALS OF BOTANY

Founded 1887

Volume 111 Number 5 May 2013

ContentSprints

i

Plant Cuttings: news in botany

v

INVITED REVIEW

- Santi, C., Bogusz, D. and Franche, C. Biological nitrogen fixation in non-legume plants 743

RESEARCH IN CONTEXT

- Foo, E., Ross, J. J., Jones, W. T. and Reid, J. B. Plant hormones in arbuscular mycorrhizal symbioses: an emerging role for gibberellins 769

ORIGINAL ARTICLES

- Budke, J. M., Goffinet, B. and Jones, C. S. Dehydration protection provided by a maternal cuticle improves offspring fitness in the moss *Funaria hygrometrica* 781

- Li, X., Bian, H., Song, D., Ma, S., Han, N., Wang, J. and Zhu, M. Flowering time control in ornamental gloxinia (*Sinningia speciosa*) by manipulation of miR159 expression 791

- de Campos, M. C. R., Pearse, S. J., Oliveira, R. S. and Lambers, H. *Viminaria juncea* does not vary its shoot phosphorus concentration and only marginally decreases its mycorrhizal colonization and cluster-root dry weight under a wide range of phosphorus supplies 801

- Kasulin, L., Agrofoglio, Y. and Botto, J. F. The receptor-like kinase *ERECTA* contributes to the shade-avoidance syndrome in a background-dependent manner 811

- El Ottra, J. H. L., Pirani, J. R. and Endress, P. K. Fusion within and between whorls of floral organs in Galipeinae (Rutaceae): structural features and evolutionary implications 821

- Jia, L., Xu, W., Li, W., Ye, N., Liu, R., Shi, L., Bin Rahman, A. N. M. R., Fan, M. and Zhang, J. Class III peroxidases are activated in proanthocyanidin-deficient *Arabidopsis thaliana* seeds 839

- Gama-Arachchige, N. S., Baskin, J. M., Geneve, R. L. and Baskin, C. C. Quantitative analysis of the thermal requirements for stepwise physical dormancy-break in seeds of the winter annual *Geranium carolinianum* (Geraniaceae) 849

- Barlow, P. W., Fisahn, J., Yazdanbakhsh, N., Moraes, T. A., Khabarova, O. V. and Gallep, C. M. *Arabidopsis thaliana* root elongation growth is sensitive to lunisolar tidal acceleration and may also be weakly correlated with geomagnetic variations 859

- Nie, Z.-L., Deng, T., Meng, Y., Sun, H. and Wen, J. Post-Boreotropical dispersals explain the pantropical disjunction in *Paederia* (Rubiaceae) 873



Contents continued on inside back cover

OXFORD OPEN

OXFORD
UNIVERSITY PRESS

Scan to view this journal
on your mobile device

Korn, R. W. The concept of the eudicot shoot apical meristem as it applies to four <i>Spiraea</i> (Rosaceae), one <i>Mentha</i> (Lamiaceae) and one <i>Euonymus</i> (Celastraceae) cultivars based on chimeric analysis	887
Ndllovu, J., Richardson, D. M., Wilson, J. R. U., O'Leary, M. and Le Roux, J. J. Elucidating the native sources of an invasive tree species, <i>Acacia pycnantha</i> , reveals unexpected native range diversity and structure	895
Miranda, T., Ebner, M., Traiser, C. and Roth-Nebelsick, A. Diurnal pattern of stomatal conductance in the large-leaved temperate liana <i>Aristolochia macrophylla</i> depends on spatial position within the leaf lamina	905
Field, D. L., Pickup, M. and Barrett, S. C. H. Ecological context and metapopulation dynamics affect sex-ratio variation among dioecious plant populations	917
Song, G. C., Choi, H. K. and Ryu, C.-M. The folate precursor para-aminobenzoic acid elicits induced resistance against <i>Cucumber mosaic virus</i> and <i>Xanthomonas axonopodis</i>	925
Pélabon, C., Osler, N. C., Diekmann, M. and Graae, B. J. Decoupled phenotypic variation between floral and vegetative traits: distinguishing between developmental and environmental correlations	935
Newton, R. J., Hay, F. R. and Ellis, R. H. Seed development and maturation in early spring-flowering <i>Galanthus nivalis</i> and <i>Narcissus pseudonarcissus</i> continues post-shedding with little evidence of maturation <i>in planta</i>	945
Quilot-Turion, B., Leppälä, J., Leinonen, P. H., Waldmann, P., Savolainen, O. and Kuittinen, H. Genetic changes in flowering and morphology in response to adaptation to a high-latitude environment in <i>Arabidopsis lyrata</i>	957
Endress, P. K., Davis, C. C. and Matthews, M. L. Advances in the floral structural characterization of the major subclades of Malpighiales, one of the largest orders of flowering plants	969
Donà, M., Balestrazzi, A., Mondoni, A., Rossi, G., Ventura, L., Buttafava, A., Macovei, A., Sabatini, M. E., Valassi, A. and Carbonera, D. DNA profiling, telomere analysis and antioxidant properties as tools for monitoring <i>ex situ</i> seed longevity	987
Purdy, S. J., Maddison, A. L., Jones, L. E., Webster, R. J., Andralojc, J., Donnison, I. and Clifton-Brown, J. Characterization of chilling-shock responses in four genotypes of <i>Miscanthus</i> reveals the superior tolerance of <i>M. × giganteus</i> compared with <i>M. sinensis</i> and <i>M. sacchariflorus</i>	999

For more information about *Annals of Botany* please visit our website at www.aob.oxfordjournals.org

Disclaimer

Statements of fact and opinion in the articles in *Annals of Botany* are those of the respective authors and contributors and not of *Annals of Botany* Company or Oxford University Press. Neither Oxford University Press nor *Annals of Botany* Company make any representation, express or implied, in respect of the accuracy of the material in this journal and cannot accept any legal responsibility or liability for any errors or omissions that may be made. The reader should make his/her own evaluation as to the appropriateness or otherwise of any experimental technique described.