



Volume 83

February 2014

ISSN 1352-2310

ATMOSPHERIC ENVIRONMENT

Air Pollution: Air Quality • Emissions • Transport and Transformations • Atmospheric Composition • Deposition and Fate • Pollution and Climate Change • Environmental Policy • Special Themes



ATMOSPHERIC ENVIRONMENT

<http://www.elsevier.com/locate/atmosenv>

Volume 83

2014

CONTENTS

Regular papers

- M. Zare Naghadehi 1 Pathological study of the prevalence of Silicosis among coal miners in Iran; A case history
- S. Candice Lung 6 Variability of Intra-urban Exposure to Particulate Matter and CO from Asian-type Community Pollution Sources
- G. Shen, M. Xue, S. Yuan, J. Zhang, Q. Zhao, B. Li, H. Wu and A. Ding 14 Chemical compositions and reconstructed light extinction coefficients of particulate matter in a mega-city in the western Yangtze River Delta, China
- N. Canha, S.M. Almeida, M.d.C. Freitas, M. Trancoso, A. Sousa, F. Mouro and H.Th. Wolterbeek 21 Particulate matter analysis in indoor environments of urban and rural primary schools using passive sampling methodology
- A. Yang, A. Jedynska, B. Hellack, I. Kooter, G. Hoek, B. Brunekreef, T.A.J. Kuhlbusch, F.R. Cassee and N.A.H. Janssen 35 Measurement of the oxidative potential of PM_{2.5} and its constituents: The effect of extraction solvent and filter type
- B.L. Zhuang, T.J. Wang, S. Li, J. Liu, R. Talbot, H.T. Mao, X.Q. Yang, C.B. Fu, C.Q. Yin, J.L. Zhu, H.Z. Che and X.Y. Zhang 43 Optical properties and radiative forcing of urban aerosols in Nanjing, China
- T. Mochizuki, A. Tani, Y. Takahashi, N. Saigusa and M. Ueyama 53 Long-term measurement of terpenoid flux above a *Larix kaempferi* forest using a relaxed eddy accumulation method
- M. Thibaudon, B. Šikoparija, G. Oliver, M. Smith and C.A. Skjøth 62 Ragweed pollen source inventory for France – The second largest centre of *Ambrosia* in Europe
- K. Mohiuddin, V. Strezov, P.F. Nelson and E. Stelcer 72 Characterisation of trace metals in atmospheric particles in the vicinity of iron and steelmaking industries in Australia
- Y. Li, F.M. Schwandner, H.J. Sewell, A. Zivkovich, M. Tigges, S. Raja, S. Holcomb, J.V. Molenaar, L. Sherman, C. Archuleta, T. Lee and J.L. Collett Jr. 80 Observations of ammonia, nitric acid, and fine particles in a rural gas production region
- M. Evtugina, C. Alves, A. Calvo, T. Nunes, L. Tarelho, M. Duarte, S.O. Prozil, D.V. Evtuguin and C. Pio 90 VOC emissions from residential combustion of Southern and mid-European woods
- M.A. Elangasinghe, K.N. Dirks, N. Singhal, S.B. Costello, I. Longley and J.A. Salmond 99 A simple semi-empirical technique for apportioning the impact of roadways on air quality in an urban neighbourhood
- X. Li, L. Feng, C. Huang, X. Yan and X. Zhang 109 Chemical characteristics of atmospheric fallout in the south of Xi'an during the dust episodes of 2001–2012 (NW China)

Continued on inside back cover

Abstracted/indexed in: *Acid Precipitation Digest, Air Pollution Titles, Analytical Abstracts, Applied Science & Technology Abstracts, Applied Science and Technology Index, Aqualine Abstracts, BIOSIS, Cambridge Scientific Abstracts, Chemical Abstracts, Current Contents / Physics / Chemistry & Earth Science, Current Contents/Agriculture, Biology & Environmental Sciences, EMBASE, Elsevier BIOBASE Environmental Periodicals Bibliography, FLUIDEX, GEOBASE, Geographical Abstracts: Physical Geography, Geological Abstracts, INSPEC, Meteorological and, Geostrophysical Abstracts, PASCAL/CNRS, Research Alert, SCISEARCH, Science Citation Index, Scopus, Theoretical Chemical Engineering Abstracts* Also covered in the abstract and citation database Scopus. Full text available on ScienceDirect®.



Printed by Polestar Wheatons Ltd., Exeter, UK



ISSN 1352-2310
83 (2014) 1–310



| | | |
|--|-----|---|
| G. de Gennaro, L. de Gennaro, A. Mazzone, F. Porcelli and M. Tutino | 119 | Indoor air quality in hair salons: Screening of volatile organic compounds and indicators based on health risk assessment |
| Y. Pang, M. Fuentes and P. Rieger | 127 | Trends in the emissions of Volatile Organic Compounds (VOCs) from light-duty gasoline vehicles tested on chassis dynamometers in Southern California |
| L.-C. Guo, L.-J. Bao, J.-W. She and E.Y. Zeng | 136 | Significance of wet deposition to removal of atmospheric particulate matter and polycyclic aromatic hydrocarbons: A case study in Guangzhou, China |
| Z. Zhang, M. Wenig, W. Zhou, T. Diehl, K.-L. Chan and L. Wang | 145 | The contribution of different aerosol sources to the Aerosol Optical Depth in Hong Kong |
| A. Verstraeten, B. De Vos, J. Neirynek, P. Roskams and M. Hens | 155 | Impact of air-borne or canopy-derived dissolved organic carbon (DOC) on forest soil solution DOC in Flanders, Belgium |
| M. Le Breton, A. Bacak, J.B.A. Muller, P. Xiao, B.M.A. Shallcross, R. Batt, M.C. Cooke, D.E. Shallcross, S.J.-B. Bauguitte and C.J. Percival | 166 | Simultaneous airborne nitric acid and formic acid measurements using a chemical ionization mass spectrometer around the UK: Analysis of primary and secondary production pathways |
| Y. Liang, H. Liang and S. Zhu | 176 | Mercury emission from coal seam fire at Wuda, Inner Mongolia, China |
| J. Sancho, J. Martínez, J.J. Pastor, J. Taboada, J.I. Piñeiro and P.J. García-Nieto | 185 | New methodology to determine air quality in urban areas based on runs rules for functional data |
| K.F. Souza, L.R.F. Carvalho, A.G. Allen and A.A. Cardoso | 193 | Diurnal and nocturnal measurements of PAH, nitro-PAH, and oxy-PAH compounds in atmospheric particulate matter of a sugar cane burning region |
| L. Mu, L. Peng, X. Liu, C. Song, H. Bai, J. Zhang, D. Hu, Q. He and F. Li | 202 | Characteristics of polycyclic aromatic hydrocarbons and their gas/particle partitioning from fugitive emissions in coke plants |
| S. Fruin, R. Urman, F. Lurmann, R. McConnell, J. Gauderman, E. Rappaport, M. Franklin, F.D. Gilliland, M. Shafer, P. Gorski and E. Avol | 211 | Spatial variation in particulate matter components over a large urban area |
| S. Yoon, S. Hu, N.Y. Kado, A. Thiruvengadam, J.F. Collins, M. Gautam, J.D. Herner and A. Ayala | 220 | Chemical and toxicological properties of emissions from CNG transit buses equipped with three-way catalysts compared to lean-burn engines and oxidation catalyst technologies |
| N. Hodas, Q. Meng, M.M. Lunden and B.J. Turpin | 229 | Toward refined estimates of ambient PM _{2.5} exposure: Evaluation of a physical outdoor-to-indoor transport model |
| S.J. White, I.M. Jamic and D.E. Angove | 237 | Chemical characterisation of semi-volatile and aerosol compounds from the photooxidation of toluene and NO _x |
| G. Zhu, X. Ma, Z. Gao, W. Ma, J. Li and Z. Cai | 245 | Characterizing CH ₄ and N ₂ O emissions from an intensive dairy operation in summer and fall in China |
| N. Lichiheb, E. Personne, C. Bedos and E. Barriuso | 260 | Adaptation of a resistive model to pesticide volatilization from plants at the field scale: Comparison with a dataset |
| R. Zhu, D. Ma and H. Xu | 269 | Summertime N ₂ O, CH ₄ and CO ₂ exchanges from a tundra marsh and an upland tundra in maritime Antarctica |
| D.H. Shin, D. Müller, T. Choi, Y.M. Noh, Y.J. Yoon, K.H. Lee, S.K. Shin, N. Chae, K. Kim and Y.J. Kim | 282 | Influence of wind speed on optical properties of aerosols in the marine boundary layer measured by ship-borne DePolarization Lidar in the coastal area of Korea |

Continued on BM IV



| | | |
|--|-----|---|
| A. Kumar | 291 | Long term (2003–2012) spatio-temporal MODIS (Terra/Aqua level 3) derived climatic variations of aerosol optical depth and cloud properties over a semi arid urban tropical region of Northern India |
| T.J. Bannan, A. Bacak, J.B.A. Muller, A.M. Booth, B. Jones, M. Le Breton, K.E. Leather, M. Ghalaieny, P. Xiao, D.E. Shallcross and C.J. Percival | 301 | Importance of direct anthropogenic emissions of formic acid measured by a chemical ionisation mass spectrometer (CIMS) during the Winter ClearfLo Campaign in London, January 2012 |
| Short communication | | |
| D.M. Moreira, A.C. Moraes, A.G. Goulart and T.T.A. Albuquerque | 254 | A contribution to solve the atmospheric diffusion equation with eddy diffusivity depending on source distance |