


TW
E58

Volume 172

January 2013

ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF


W.J. Manning

ASSOCIATE EDITORS

B. Nowack

C. Wiegand

E. Paoletti



CONTENTS

Volume 172 January 2013

- 1 Gonadal abnormalities in frogs (*Lithobates* spp.) collected from managed wetlands in an agricultural region of Nebraska, USA**
D.M. Papoulias, M.S. Schwarz, L. Mena
Frogs from wetlands in agricultural areas had reproductive anomalies however no statistical relationship existed between biomarkers and measured concentrations of pesticides.
- 9 Chemical stabilization of metals and arsenic in contaminated soils using oxides – A review**
M. Komárek, A. Vaněk, V. Ettler
In situ stabilization of metals and As in contaminated soils using oxides combined with phytostabilization is a potential alternative to conventional remediation techniques.
- 23 Toxic effects of erythromycin, ciprofloxacin and sulfamethoxazole exposure to the antioxidant system in *Pseudokirchneriella subcapitata***
X.-P. Nie, B.-Y. Liu, H.-J. Yu, W.-Q. Liu, Y.-F. Yang
Antibiotics (Erythromycin, ciprofloxacin and sulfamethoxazole) cause the change of antioxidant system and lead to oxidative stress to a green microalga, Pseudokirchneriella subcapitata.
- 33 Bioaugmentation with a consortium of bacterial nitrophenol-degraders for remediation of soil contaminated with three nitrophenol isomers**
X.-Q. Chi, J.-J. Zhang, S. Zhao, N.-Y. Zhou
A soil contaminated with three nitrophenol isomers has been successfully bioremediated by inoculation with a bacterial consortium.
- 42 Intra- and inter-basin mercury comparisons: Importance of basin scale and time-weighted methylmercury estimates**
P.M. Bradley, C.A. Journey, M.E. Brigham, D.A. Burns, D.T. Button, K. Riva-Murray
Fluvial methylmercury concentration correlates with wetland area not basin scale and time-weighted estimates better predict basin top predator mercury than discrete sample estimates.
- 53 Changes in Sb speciation with waterlogging of shooting range soils and impacts on plant uptake**
X.-m. Wan, S. Tandy, K. Hockmann, R. Schulin
Antimony uptake by grasses was dependant on redox speciation but responses differed between species.
- 61 Biomonitoring of urban air pollution: Magnetic studies and SEM observations of corticolous foliose and microfoliose lichens and their suitability for magnetic monitoring**
M.A.E. Chaparro, J.M. Lavernia, M.A.E. Chaparro, A.M. Sinito
Lichen species are airborne pollutant collectors that are well-distributed in some urban areas and allow effective magnetic monitoring at low cost in cities.
- 70 Effects of selenium accumulation on phytotoxicity, herbivory, and pollination ecology in radish (*Raphanus sativus* L.)**
K.R. Hladun, D.R. Parker, K.D. Tran, J.T. Trumble
*Radish accumulated the pollutant selenium in floral tissues, but this did not deter the pollinator (*Apis mellifera*) from foraging.*
- 76 Ecotoxicity of manufactured ZnO nanoparticles – A review**
H. Ma, P.L. Williams, S.A. Diamond
Ecotoxicity of ZnO nanoparticles.
- 86 Enhanced PCBs sorption on biochars as affected by environmental factors: Humic acid and metal cations**
Y. Wang, L. Wang, G. Fang, H.M.S.K. Herath, Y. Wang, L. Cang, Z. Xie, D. Zhou
The biochars had higher adsorption affinity for PCBs in the extracted soil solution because coexisting humic acid and metal cations increased their sorption.

Continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information-Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; Thomson Scientific GeoSciTech; Science Citation Index; SciSearch). Also covered in the abstract and citation database SciVerse Scopus®. Full text available on SciVerse ScienceDirect®.



0269-7491(201301)172;1-Q



ELSEVIER

ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 94 Primary investigation on contamination pattern of legacy and emerging halogenated organic pollutions in freshwater fish from Liaohe River, Northeast China**
G. Ren, Z. Wang, Z. Yu, Y. Wang, S. Ma, M. Wu, G. Sheng, J. Fu
An unusually high percentage of PCB-209 was first reported in the fish samples collected from China, which might indicate that there were distinct sources of pure PCB-209 in the region of Liaohe River.
- 100 Fluorescence quenching of fulvic acids by fullerene in water**
F. Wu, Y. Bai, Y. Mu, B. Pan, B. Xing, Y. Lin
Interactions between nanoscaled-fullerene-aggregates and fulvic acids in water were quantitatively investigated by fluorescence quenching titration.
- 108 Monitoring bioremediation of atrazine in soil microcosms using molecular tools**
S. Sagarkar, S. Mukherjee, A. Nousiainen, K. Björklöf, H.J. Purohit, K.S. Jørgensen, A. Kapley
This study demonstrates the use of molecular tools in bioremediation to monitor and track target genes; correlates the results with degradation and thereby predicts the efficiency of treatment.
- 116 Incidence of invasive macrophytes on methylmercury budget in temperate lakes: Central role of bacterial periphytic communities**
S. Gentès, M. Monperrus, A. Legeay, R. Maury-Brachet, S. Davail, J.-M. André, R. Guyoneaud
The incidence of periphytic microbial communities on net methylmercury production is highlighted in temperate aquatic ecosystems.
- 124 Migration of polybrominated diphenyl ethers in biosolids-amended soil**
T. Gorgy, L.Y. Li, J.R. Grace, M.G. Ikononou
PBDEs migrated downward over one year after applying biosolids to agricultural soil, but photodegradation or volatilization appear to have also occurred.
- 131 Spatial scales of pollution from variable resolution satellite imaging**
A.A. Chudnovsky, A. Kostinski, A. Lyapustin, P. Koutrakis
The correlation between $PM_{2.5}$ and AOD decreases as AOD resolution is degraded. The new high-resolution MAIAC AOD retrieval has the potential to capture $PM_{2.5}$ variability at the intra-urban scale.
- 139 Comparing early twentieth century and present-day atmospheric pollution in SW France: A story of lichens**
Y. Agnan, N. Séjalon-Delmas, A. Probst
Using lichens registration, past (As, Pb, Cd) and recent (Sb, Sn, Cu) atmospheric pollution from local and long-range transport, as well as dust deposition, was evidenced in SW France.
- 149 Arsenic speciation in Chinese Herbal Medicines and human health implication for inorganic arsenic**
X.-J. Liu, Q.-L. Zhao, G.-X. Sun, P. Williams, X.-J. Lu, J.-Z. Cai, W.-J. Liu
Inorganic arsenic was the predominant species in all of CHMs samples.
- 155 Integrated monitoring of particle associated transport of PAHs in contrasting catchments**
M. Schwientek, H. Rügner, B. Beckingham, B. Kuch, P. Grathwohl
Particle-facilitated transport of PAHs was found to relate to urban population pressure relative to suspended particle loading in contrasting catchments.
- 163 Assessment of flame retardants in river water using a ceramic dosimeter passive sampler**
J. Cristale, A. Katsoyiannis, C. Chen, K.C. Jones, S. Lacorte
Ceramic dosimeter using HLB as sorbent can be an efficient river-water passive sampler for organophosphorous and brominated flame retardants.
- 170 Maximum entropy estimation of a Benzene contaminated plume using ecotoxicological assays**
A. Wahyudi, M. Bartzke, E. Küster, P. Bogaert
A Maximum Entropy method to rebuild qualitative relationships between Benzene groundwater concentrations and their ecotoxicological effect.
- 180 Short-term extractability of sulfadiazine after application to soils**
T. Müller, I. Rosendahl, A. Focks, J. Siemens, J. Klasmeier, M. Matthies
The extractability of sulfadiazine from soil is reduced by an instantaneous process on a time scale of minutes followed by a kinetic sequestration on a time scale of days.
- 186 Metal exposure and accumulation patterns in free-range cows (*Bos taurus*) in a contaminated natural area: Influence of spatial and social behavior**
S. Roggeman, N. van den Brink, N. Van Praet, R. Blust, L. Bervoets
This study reveals that spatial heterogeneity and foraging behavior play a more important role in the metal exposure pattern of large grazers than generally is presumed.
- 200 Life-history effects of arsenic toxicity in clades of the earthworm *Lumbricus rubellus***
C.J. Anderson, P. Kille, A.J. Lawlor, D.J. Spurgeon
*The impact of arsenic toxicity and role of genotype on *Lumbricus rubellus* life-history parameters reveal variation among sublethal endpoints.*

CONTENTS—Continued from inside back cover

- 208 Identification and dynamic modeling of biomarkers for bacterial uptake and effect of sulfonamide antimicrobials**
M.K. Richter, A. Focks, B. Siegfried, D. Rentsch, M. Krauss, R.P. Schwarzenbach, J. Hollender
Identification of specific biomarkers for the uptake and effect-based reaction of sulfonamides in bacteria and resulting growth inhibition.
- 216 Mercury trends in herring gull (*Larus argentatus*) eggs from Atlantic Canada, 1972–2008: Temporal change or dietary shift?**
N.M. Burgess, A.L. Bond, C.E. Hebert, E. Neugebauer, L. Champoux
Mercury levels have remained stable in herring gull eggs at four of five breeding colonies monitored in Atlantic Canada from the 1970s to 2008.
- 223 Methods to assess bioavailability of hydrophobic organic contaminants: Principles, operations, and limitations**
X. Cui, P. Mayer, J. Gan
This review summarizes the principles and operations of bioavailability prediction methods, discusses their strengths and limitations, and highlights issues for future research.
- 235 Trophodynamics of inorganic pollutants in a wide-range feeder: The relevance of dietary inputs and biomagnification in the Yellow-legged gull (*Larus michahellis*)**
R. Ramos, F. Ramirez, L. Jover
Pollutant body burden of a wide-range feeder was highly influenced by dietary exposure (using $\delta^{34}\text{S}$ as proxy), whereas the influence of trophic position (using $\delta^{15}\text{N}$) was relatively low.
- 243 Physiological aspects of mangrove (*Laguncularia racemosa*) grown in microcosms with oil-degrading bacteria and oil contaminated sediment**
V. Sodr , V.S. Caetano, R.M. Rocha, F.L. Carmo, L.O. Medici, R.S. Peixoto, A.S. Rosado, F. Reinert
Despite the need to establish methods to diagnose the health status of mangroves little is known about the impacts of petrochemicals on mangrove plants and associated rhizosphere microorganisms.
- 250 Assessing ozone and nitrogen impact on net primary productivity with a Generalised non-Linear Model**
A. De Marco, A. Screpanti, F. Attorre, C. Proietti, M. Vitale
Ozone and nitrogen depositions have non-linear effects on primary productivity of tree species differently distributed in Italy.
- 264 The ecological effects of a herbicide–insecticide mixture on an experimental freshwater ecosystem**
C.B. Choung, R.V. Hyne, M.M. Stevens, G.C. Hose
A low concentration of terbufos results in delayed population level effects.
- 275 Effects of chlorimuron ethyl on terrestrial and wetland plants: Levels of, and time to recovery following sublethal exposure**
D. Carpenter, C. Boutin, J.E. Allison
This study serves to bridge the gap between simplified short-term greenhouse tests and effects of herbicides on recovery of non-target plant species after sublethal exposures.