


ПН
E58

Volume 173

February 2013

ISSN 0269-7491




ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

**B. Nowack
C. Wiegand
E. Paoletti**



CONTENTS

Volume 173 February 2013

- 1 Mapping urban pipeline leaks: Methane leaks across Boston**
N.G. Phillips, R. Ackley, E.R. Crosson, A. Down, L.R. Hutrya, M. Brondfield, J.D. Karr, K. Zhao, R.B. Jackson
We identified 3356 methane leaks in Boston, with isotopic characteristics consistent with pipeline natural gas.
- 5 Synergistic toxicity of Macondo crude oil and dispersant Corexit 9500A® to the *Brachionus plicatilis* species complex (Rotifera)**
R. Rico-Martínez, T.W. Snell, T.L. Shearer
*Using *Brachionus plicatilis* acute toxicity tests we estimated Corexit 9500A® and oil toxicity. When these compounds are mixed a 52-fold increase in toxicity was observed.*
- 11 Influence of contact time and sediment composition on the bioavailability of Cd in sediments**
H. Zhong, L. Kraemer, D. Evans
Transfer of Cd from carbonate phase to Fe–Mn oxide phase in sediments was important in affecting the decrease of Cd bioavailability over time.
- 17 Biodegradation of 17 α -ethinylestradiol by heterotrophic bacteria**
S. Larcher, V. Yargeau
*Heterotrophic bacteria, especially *Rhodococcus* species, are capable of successfully degrading 17 α -ethinylestradiol (EE2).*
- 23 Responses of soil Collembola to long-term atmospheric CO₂ enrichment in a mature temperate forest**
G.-L. Xu, S.-L. Fu, P. Schleppei, M.-H. Li
Seven years of CO₂ enrichment caused a significant decrease in Collembola abundance, especially in euedaphic species.
- 29 Upward and downward solar-induced chlorophyll fluorescence yield indices of four tree species as indicators of traffic pollution in Valencia**
S. Van Wittenbergh, L. Alonso, J. Verelst, I. Hermans, J. Delegido, F. Veroustraete, R. Valcke, J. Moreno, R. Samson
Fluorescence yield parameters as stress detectors for traffic-exposed urban trees are tested.
- 38 Nanoscale zerovalent iron alters soil bacterial community structure and inhibits chloroaromatic biodegradation potential in Aroclor 1242-contaminated soil**
E.L. Tilston, C.D. Collins, G.R. Mitchell, J. Princivalle, L.J. Shaw
nZVI inhibits microbial functions of potential importance for remediation strategies combining nZVI treatment and biodegradation.
- 47 Using SPME fibers and Tenax to predict the bioavailability of pyrethroids and chlorpyrifos in field sediments**
A.D. Harwood, P.F. Landrum, D.P. Weston, M.J. Lydy
This research provides an important first step in using bioavailability-based techniques for estimating the bioavailability and toxicity of hydrophobic pesticides in field sediments.
- 52 Recent changes in mercury deposition and primary productivity inferred from sediments of lakes from the Hudson Bay Lowlands, Ontario, Canada**
M.L. Brazeau, A.J. Poulain, A.M. Paterson, W.(Bill) Keller, H. Sanei, J.M. Blais
Increased mercury concentrations in lake sediment cores coincide with evidence of increased autochthonous production in lakes of the Hudson Bay Lowlands, Canada.
- 61 Trophic transfer of pyrene metabolites between aquatic invertebrates**
V. Carrasco Navarro, M.T. Leppänen, J.V.K. Kukkonen, S. Godoy Olmos
*Some of the pyrene metabolites produced by the model invertebrates *L. variegatus* and *C. riparius* are transferred to *G. setosus* through the diet, proving their trophic transfer.*

Continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information-Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; Thomson Scientific GeoSciTech; Science Citation Index; SciSearch). Also covered in the abstract and citation database SciVerse Scopus®. Full text available on SciVerse ScienceDirect®.



0269-7491(201302)173;1-0



ELSEVIER



CONTENTS—Continued from outside back cover

- 68 Pb and Zn imaging and *in situ* speciation at the geogenic/biogenic interface in sentinel earthworms using electron microprobe and synchrotron micro-focus X-ray spectroscopy**
A.J. Morgan, P. Kille, A. Bennett, M. O'Reilly, P. Fisher, J.M. Charnock
An in situ study of metal imaging and speciation in an environment model, the earthworm, using an innovative combination of synchrotron and electron probe μ -focus X-ray analyses.
- 75 Ecotoxicological effects evoked in hydrophytes by leachates of invasive *Acer negundo* and autochthonous *Alnus glutinosa* fallen off leaves during their microbial decomposition**
A. Krevš, J. Darginavičienė, B. Gylytė, R. Grigutytė, S. Jurkonienė, R. Karitonas, A. Kučinskienė, R. Pakalnis, K. Sadauskas, R. Vitkus, L. Manusadžianas
Fallen leaves and leaf litter leachates from invasive and native species of trees, which enter waterbody can be environmental factor affecting differently microbial biodestruction and aquatic vegetation in freshwater systems, thus influencing ecological scenarios.
- 85 Effects of ozone on stomatal responses to environmental parameters (blue light, red light, CO₂ and vapour pressure deficit) in three *Populus deltoides* × *Populus nigra* genotypes**
J. Dumont, F. Spicher, P. Montpied, P. Dizengremel, Y. Jolivet, D. Le Thiec
Taking into account stomatal behaviour upon exposure to high ozone levels is essential for improving phytotoxic ozone dose assessment.
- 97 Effects of water chemistry on the dissolution of ZnO nanoparticles and their toxicity to *Escherichia coli***
M. Li, D. Lin, L. Zhu
The toxicity of nano-ZnO to E. coli depended on not only free Zn²⁺ but also Ca²⁺ and Mg²⁺ which could reduce the toxicity of Zn²⁺.
- 103 High altitude artisanal small-scale gold mines are hot spots for Mercury in soils and plants**
T.A. Terán-Mita, A. Faz, F. Salvador, J.M. Arocena, J.A. Acosta
Mean contents of Hg in soils were at least 5× to 60× more compared to Hg in control sites, and the high Hg in Poaceae and Rosaceae may elevate Hg into the food chain.
- 110 A spatially-based modeling framework for assessing the risks of soil-associated metals to bats**
B.V. Hernout, K.E. Somerwill, K.E. Arnold, C.J. McClean, A.B.A. Boxall
Application of a novel, spatially explicit risk assessment framework indicates that the health of insectivorous bat species in some regions of the UK may be at threat from exposure to soil associated metals.
- 117 Metal concentrations in stream biofilm and sediments and their potential to explain biofilm microbial community structure**
P.-Y. Ancion, G. Lear, A. Dopheide, G.D. Lewis
Metal concentrations in stream biofilms provide a good assessment of the effects of trace metal contaminants on freshwater ecosystems.
- 125 Biofuel components change the ecology of bacterial volatile petroleum hydrocarbon degradation in aerobic sandy soil**
A. Elazhari-Ali, A.K. Singh, R.J. Davenport, I.M. Head, D. Werner
Blending of petroleum with ethanol or biodiesel changes the fuel degrading soil bacterial community structure, but the long-term effects on fuel biodegradability are minor.
- 133 Occurrence and dynamics of micropollutants in a karst aquifer**
B. Morasch
Regional UPLC-MS/MS-based survey underlines the vulnerability of a Swiss karst system towards micropollutants and the potential exposure to atrazine through karst drinking water.
- 138 Evaluating nutrient impacts in urban watersheds: Challenges and research opportunities**
R.O. Carey, G.J. Hochmuth, C.J. Martinez, T.H. Boyer, M.D. Dukes, G.S. Toor, J.L. Cisar
Urban watersheds include numerous nitrogen and phosphorus sources; some sources have been quantified, but several opportunities still exist to evaluate nutrients in urban watersheds.
- 150 Pollution level, inhalation exposure and lung cancer risk of ambient atmospheric polycyclic aromatic hydrocarbons (PAHs) in Taiyuan, China**
Z. Xia, X. Duan, S. Tao, W. Qiu, D. Liu, Y. Wang, S. Wei, B. Wang, Q. Jiang, B. Lu, Y. Song, X. Hu
The inhalation exposure and lung cancer risk of ambient atmospheric PAHs changed for different seasons, areas and population groups in Taiyuan, China.
- 157 Salinisation of rivers: An urgent ecological issue**
M. Cañedo-Argüelles, B.J. Kefford, C. Piscart, N. Prat, R.B. Schäfer, C.-J. Schulz
Secondary salinization of the world rivers and streams.
- 168 Enhanced degradation of ¹⁴C-HCB in two tropical clay soils using multiple anaerobic-aerobic cycles**
F.O. Kengara, U. Doerfler, G. Welzl, B. Ruth, J.C. Munch, R. Schroll
Two anaerobic-aerobic cycles enhance the degradation of HCB in two ecologically different tropical clay soils.
- 176 Dechlorane Plus and its dechlorinated analogs from an e-waste recycling center in maternal serum and breast milk of women in Wenling, China**
Y.-J. Ben, X.-H. Li, Y.-L. Yang, L. Li, J.-P. Di, W.-Y. Wang, R.-F. Zhou, K. Xiao, M.-Y. Zheng, Y. Tian, X.-B. Xu
DP can bio-accumulate in human with the low milk/serum partition coefficient and similar blood and milk stereo-selective bio-accumulation profiles.

- 182 Beyond the bed: Effects of metal contamination on recruitment to bedded sediments and overlying substrata**
N.A. Hill, S.L. Simpson, E.L. Johnston
Under natural disturbance regimes, metal-contaminated sediments pose less of a direct risk to hard-substratum fauna than to sediment-dwelling fauna and SQG appear appropriate.
- 192 Comparison of extended-spectrum- β -lactamase (ESBL) carrying *Escherichia coli* from sewage sludge and human urinary tract infection**
G. Zarfel, H. Galler, G. Feierl, D. Haas, C. Kittinger, E. Leitner, A.J. Grisold, F. Mascher, J. Posch, B. Pertschy, E. Marth, F.F. Reinthaler
The distribution of ESBL resistance genes in isolates from patients and environmental samples.
- 200 Does selective serotonin reuptake inhibitor (SSRI) fluoxetine affects mussel *Mytilus galloprovincialis*?**
M. Gonzalez-Rey, M.J. Bebianno
*Exposure to 75 ng L⁻¹ antidepressant fluoxetine (FLX) induces tissue-specific multibiomarker responses alteration in mussel *Mytilus galloprovincialis*.*
- 210 In vitro assessment of thyroid hormone disrupting activities in drinking water sources along the Yangtze River**
X. Hu, W. Shi, F. Zhang, F. Cao, G. Hu, Y. Hao, S. Wei, X. Wang, H. Yu
Drinking water sources from the lower reaches of Yangtze River showed thyroid hormone disrupting potential in reporter gene assays.
- 216 Estimation of the algal-available phosphorus pool in sediments of a large, shallow eutrophic lake (Taihu, China) using profiled SMT fractional analysis**
M. Zhu, G. Zhu, W. Li, Y. Zhang, L. Zhao, Z. Gu
Profiled SMT fractional analysis of internal phosphorus pool in large, shallow lake.
- 224 Distribution of hydrocarbons released during the 2010 MC252 oil spill in deep offshore waters**
C. Spier, W.T. Stringfellow, T.C. Hazen, M. Conrad
All available certified Deepwater Horizon data was used to determine the spatial, temporal, and chemical distribution of hydrocarbons in subsurface of the Gulf of Mexico.
- 231 Effects of plant species identity, diversity and soil fertility on biodegradation of phenanthrene in soil**
A.O. Oyelami, U.V. Okere, K.H. Orwin, G.B. De Deyn, K.C. Jones, K.T. Semple
Soil management was the main driver for the mineralisation of ¹⁴C-phenanthrene in soil.
- 238 Arsenic and selenium mobilisation from organic matter treated mine spoil with and without inorganic fertilisation**
E. Moreno-Jiménez, R. Clemente, A. Mestrot, A.A. Meharg
The addition of organic amendment and/or inorganic fertiliser to a trace element contaminated soil in flooded conditions led to As, Cu and Se solubilisation.
- 245 Terrestrial mosses as biomonitors of atmospheric POPs pollution: A review**
H. Harmens, L. Foan, V. Simon, G. Mills
Mosses are suitable biomonitors of persistent organic pollutants (POPs).
- 255 Air Quality Standards for Particulate Matter (PM) at high altitude cities**
H. Bravo Alvarez, R. Sosa Echeverria, P. Sanchez Alvarez, S. Krupa
In order to compare high altitude atmospheric PM concentrations with AQS, one must consider T and P of the sampling site.
- 257 An endocrine disruptor, bisphenol A, affects development in the protochordate *Ciona intestinalis*: Hatching rates and swimming behavior alter in a dose-dependent manner**
A. Matsushima, K. Ryan, Y. Shimohigashi, I.A. Meinertzhagen
*Exposure of fertilized *Ciona* embryos to BPA decreased their hatch rate in a dose-dependent manner and led to abnormal larval swimming behavior.*
- 264 Increased frequency and severity of developmental deformities in rough-skinned newt (*Taricha granulosa*) embryos exposed to road deicing salts (NaCl & MgCl₂)**
G.R. Hopkins, S.S. French, E.D. Brodié Jr.
Two commonly used road deicing salts, NaCl and MgCl₂, caused increased frequency and severity of developmental deformities in rough-skinned newt embryos.
- 270 Assessment of potential climate change impacts on peatland dissolved organic carbon release and drinking water treatment from laboratory experiments**
R. Tang, J.M. Clark, T. Bond, N. Graham, D. Hughes, C. Freeman
Future drought events are likely to alter soil moisture, which predominately controls production of peat-derived dissolved organic carbon and subsequently drinking water quality.