

774
E58

Volume 177

June 2013

ISSN 0269-7491



ELSEVIER

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

J. Gan

B. Nowack

E. Paoletti

C. Wiegand



CONTENTS

Volume 177 June 2013

- 1 Trophic level transfer of microplastic: *Mytilus edulis* (L.) to *Carcinus maenas* (L.)**
P. Farrell, K. Nelson
This communication demonstrates trophic level transfer of microplastic particles from Mytilus edulis to Carcinus maenas.
- 4 CO₂ absorption/emission and aerodynamic effects of trees on the concentrations in a street canyon in Guangzhou, China**
J.-F. Li, J.-M. Zhan, Y.S. Li, O.W.H. Wai
The effects of trees on CO₂ concentrations in a street canyon are examined by CFD simulations, taking into account both the CO₂ absorption/emission and aerodynamic effects of trees.
- 13 A three-dimensional water quality modeling approach for exploring the eutrophication responses to load reduction scenarios in Lake Yilong (China)**
L. Zhao, Y. Li, R. Zou, B. He, X. Zhu, Y. Liu, J. Wang, Y. Zhu
The lake water quality responds to watershed load reduction in a nonlinear way, which requires combined watershed and in-lake management for lake restoration.
- 22 Effects of naphthenic acid exposure on development and liver metabolic processes in anuran tadpoles**
S.D. Melvin, C.M. Lanctôt, P.M. Craig, T.W. Moon, K.M. Peru, J.V. Headley, V.L. Trudeau
*Leopard frog (*Lithobates pipiens*) tadpoles chronically exposed to sub-lethal NA concentrations (1–2 mg/L) suffered decreased growth and development and disruption to liver metabolic processes.*
- 28 Distributions and sea-to-air fluxes of chloroform, trichloroethylene, tetrachloroethylene, chlorodibromomethane and bromoform in the Yellow Sea and the East China Sea during spring**
Z. He, G.-P. Yang, X.-L. Lu, H.-H. Zhang
The elevated levels of the volatile halocarbons in the coastal waters were attributed to anthropogenic inputs and biological release by phytoplankton (e.g., diatoms).
- 38 Effect of organic matter amendment, arsenic amendment and water management regime on rice grain arsenic species**
G.J. Norton, E.E. Adomako, C.M. Deacon, A.-M. Carey, A.H. Price, A.A. Meharg
The addition of high amounts of organic matter to soils led to an increase in total rice grain arsenic, as well as alteration in the percentage arsenic species in the rice grains.
- 48 A novel *in situ* method for sampling urban soil dust: Particle size distribution, trace metal concentrations, and stable lead isotopes**
X. Bi, S. Liang, X. Li
*Soil dust collected by a novel *in situ* sampling method can provide critical information about contamination and potential human exposure in urban environments.*
- 58 Passive sampling of perfluorinated chemicals in water: Flow rate effects on chemical uptake**
S.L. Kaserzon, E.L.M. Vermeirssen, D.W. Hawker, K. Kennedy, C. Bentley, J. Thompson, K. Booij, J.F. Mueller
A modified polar organic chemical integrative sampler (POCIS) shows promise as a sampling tool to support monitoring efforts for PFCs in a range of water flow rate conditions.
- 64 Distribution and speciation of metals (Cu, Zn, Cd, and Pb) in agricultural and non-agricultural soils near a stream upriver from the Pearl River, China**
S. Yang, D. Zhou, H. Yu, R. Wei, B. Pan
The mobility of four investigated metals is low but Cd pollution should not be overlooked in soils upstream of Pearl River.
- 71 Hydroxylated PBDEs and brominated phenolic compounds in particulate matters emitted during recycling of waste printed circuit boards in a typical e-waste workshop of South China**
Z. Ren, X. Bi, B. Huang, M. Liu, G. Sheng, J. Fu
Recycling of waste printed circuit boards might represent a strong source of OH-PBDEs, BPs and other phenolic compounds.

Continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information-Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; Thomson Scientific GeoSciTech; Science Citation Index; SciSearch). Also covered in the abstract and citation database SciVerse Scopus®. Full text available on SciVerse ScienceDirect®.



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 78 The effects of aluminum and nickel in nectar on the foraging behavior of bumblebees**
G.A. Meindl, T.-L. Ashman
We extend current understanding of the effects of plant chemistry on plant-pollinator interactions by describing the effects of metals in nectar on bee foraging.
- 82 Biomonitoring study of an estuarine coastal ecosystem, the Sacca di Goro lagoon, using *Ruditapes philippinarum* (Mollusca: Bivalvia)**
A. Sacchi, C. Mouneyrac, C. Bolognesi, A. Sciutto, P. Roggieri, M. Fusi, G.M. Beone, E. Capri
*The bivalve *Ruditapes philippinarum* as a bioindicator in monitoring pollution of estuaries.*
- 90 Extractability and bioavailability of Pb and As in historically contaminated orchard soil: Effects of compost amendments**
M. Fleming, Y. Tai, P. Zhuang, M.B. McBride
Remediating metal-contaminated orchard soils with compost reduced lead bioavailability but had little effect on arsenic.
- 98 Experimental increase in availability of a PAH complex organic contamination from an aged contaminated soil: Consequences on biodegradation**
A. Cébron, P. Faure, C. Lorgeoux, S. Ouvrard, C. Leyval
More than ageing, the complexity of organic contamination is the main driver of PAH availability.
- 106 Cumulative impacts of human activities on urban garden soils: Origin and accumulation of metals**
Zs. Szolnoki, A. Farsang, I. Puskás
Metal enrichment and sources in urban garden soils due to urban activities and cultivation were successfully identified by combining more methods (enrichment factors, statistical analysis, spatial distribution).
- 116 Atmospheric concentrations and potential sources of PCBs, PBDEs, and pesticides to Acadia National Park**
S.C. Sofuoglu, A. Sofuoglu, T.M. Holsen, C.M. Alexander, J.J. Pagano
This study concludes that the high-pollutant concentrations arriving at ANP do not exclusively originate from the major urban centers along the eastern Atlantic seaboard.
- 125 Building statistical associations to forecast ethylbenzene levels in European urban-traffic environments**
Ch. Vlachokostas, A.V. Michailidou, D. Spyridi, N. Moussiopoulos
Multi-stressor statistical analysis is capable of capturing ethylbenzene concentration trends in EU urban environments and could support decision-making.
- 135 Within- and among-population level differences in response to chronic copper exposure in southern toads, *Anaxyrus terrestris***
S.L. Lance, R.W. Flynn, M.R. Erickson, D.E. Scott
CapsuleSouthern toad aquatic survival is significantly reduced at 15 µg Cu/L with variation in survivorship across clutches and populations.
- 143 Exposure to a widespread non-pathogenic bacterium magnifies sublethal pesticide effects in the damselfly *Enallagma cyathigerum*: From the suborganismal level to fitness-related traits**
L. Janssens, R. Stoks
Non-pathogenic bacteria reduce fitness-related traits and can synergistically interact with sublethal pesticide effects for physiological and fitness-related traits.
- 150 Sex-dependent accumulation and maternal transfer of Dechlorane Plus flame retardant in fish from an electronic waste recycling site in South China**
J.-P. Wu, Y.-Z. She, Y. Zhang, Y. Peng, L. Mo, X.-J. Luo, B.-X. Mai
Sex-dependent accumulation and species-specific maternal transfer of Dechlorane Plus flame retardant was observed in two highly contaminated fish species.
- 156 Prediction of N₂O emission from local information with Random Forest**
A. Philibert, C. Loyce, D. Makowski
Random Forest, a machine learning method, outperformed the regression models for predicting N₂O emissions and led to the identification of three important input variables.
- 164 Non-destructive techniques for biomonitoring of spatial, temporal, and demographic patterns of mercury bioaccumulation and maternal transfer in turtles**
B.C. Hopkins, M.J. Hepner, W.A. Hopkins
We developed and validated mathematical models that will be useful for biomonitoring Hg accumulation in turtles in a conservation-minded fashion.
- 171 Key factors affecting liver PBDE concentrations in sparrowhawks (*Accipiter nisus*)**
J.D. Crosse, R.F. Shore, K.C. Jones, M.G. Pereira
Liver PBDE concentrations in UK sparrowhawks are amongst the highest reported anywhere but vary markedly with body condition and age class.
- 177 Comparing ozone metrics on associations with outpatient visits for respiratory diseases in Taipei Metropolitan area**
Y.-K. Lin, S.-C. Chang, C. Lin, Y.-C. Chen, Y.-C. Wang
The O_{3, 24 h avg} was a better metric than O_{3, 1 h max} and O_{3, 8 h max} to associate with outpatient visits of total respiratory diseases.

CONTENTS—Continued from inside back cover

185 Toxicity testing of dispersed oil requires adherence to standardized protocols to assess potential real world effects

G. Coelho, J. Clark, D. Aurand

To effectively use toxicity testing data from lab tests with oils and chemically dispersed oils, researchers must follow standardized testing protocols.

189 Impacts of increasing ozone on Indian plants

E. Oksanen, V. Pandey, A.K. Pandey, S. Keski-Saari, S. Kontunen-Soppela, C. Sharma

Tropospheric ozone is an increasing threat to food production in India.

Letter to the Editor

201 Implications of greater than average increases in nitrogen deposition on the western edge of the Szechwan Basin, China

Z. Xu, L. Tu, T. Hu, M. Schädler

Reply to Letter to the Editor

203 Reply to “Implications of greater than average increases in nitrogen deposition on the western edge of the Szechwan Basin, China” by Xu et al. (2013)

X. Liu

Response

204 Response to authors’ reply regarding “Implications of greater than average increases in nitrogen deposition on the western edge of the Szechwan Basin, China” by Xu et al. (2013)

Z. Xu