


711  
E58

Volume 180

September 2013

ISSN 0269-7491



# ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

**W.J. Manning**


ASSOCIATE EDITORS

**J. Gan**

**B. Nowack**

**E. Paoletti**

**C. Wiegand**




## CONTENTS

### Volume 180 September 2013

- 1 Sorption of humic acid to functionalized multi-walled carbon nanotubes**  
F. Wang, J. Yao, H. Chen, Z. Yi, B. Xing  
*The functionalities of MCNTs regulated the sorption behavior between MCNTs and HA.*
- 7 Fate and movement of selenium from drainage sediments disposed onto soil with and without vegetation**  
G.S. Bañuelos, C. Bitterli, R. Schulin  
*Phytomanagement was not effective in preventing the downward migration of selenium deposited as drainage sediment on light textured soil.*
- 13 Trend of dissolved inorganic nitrogen at stations downstream from the Three-Gorges Dam of Yangtze River**  
C.C. Sun, Z.Y. Shen, M. Xiong, F.B. Ma, Y.Y. Li, L. Chen, R.M. Liu  
*DIN concentrations were reflections of agricultural development in the YRB, but the impoundment of TGD since June, 2003 also had impacts on DIN concentrations far to the Hankou station.*
- 19 Assessment of PCDD/Fs levels in soil at a contaminated sawmill site in Sweden – A GIS and PCA approach to interpret the contamination pattern and distribution**  
S. Henriksson, J. Hagberg, M. Bäckström, I. Persson, G. Lindström  
*The use of geostatistical and multivariate statistical methods are powerful tools to visualize the contamination pattern and distribution at a highly PCDD/Fs-contaminated site.*
- 27 Spatial distribution of polychlorinated naphthalenes in the atmosphere across North China based on gridded field observations**  
Y. Lin, Y. Zhao, X. Qiu, J. Ma, Q. Yang, M. Shao, T. Zhu  
*High levels of PCNs were observed in the atmosphere of North China, which were mainly from thermal processes.*
- 34 Evaluating the sub-lethal toxicity of PFOS and PFOA using rotifer *Brachionus calyciflorus***  
L. Zhang, J. Niu, Y. Li, Y. Wang, D. Sun  
*The toxicity data of PFOS and PFOA on the rotifer, at both individual and population levels, can enhance the potential of rotifers as suitable aquatic models for ecotoxicology tests.*
- 41 New insights into parental effects and toxicity: Mate availability and diet in the parental environment affect offspring responses to contaminants**  
S.C. Plautz, M.A. Funkhouser, C.J. Salice  
*Both parental diet composition and mating strategy can significantly alter the toxicant tolerance of offspring, and toxicant characteristics likely influence the probability of parental effects.*
- 48 Covalent bonding of chloroanilines to humic constituents: Pathways, kinetics, and stability**  
D. Kong, Q. Xia, G. Liu, Q. Huang, J. Lu  
*Bound residues of chloroanilines formed via 1,2- and 1,4-nucleophilic addition pathways showed different stability.*
- 55 Responses of ecosystem carbon dioxide exchange to nitrogen addition in a freshwater marshland in Sanjiang Plain, Northeast China**  
L. Zhang, C. Song, P.N. Nkrumah  
*Effects of increases in nitrogen deposition from anthropogenic sources on carbon sequestration.*
- 63 Ecotoxicity of non-aged and aged CeO<sub>2</sub> nanomaterials towards freshwater microalgae**  
N. Manier, A. Bado-Nilles, P. Delalain, O. Aguerre-Chariol, P. Pandard  
*Does agglomeration state influence the ecotoxicity of engineered nanomaterials?*
- 71 Subacute toxicity of copper and glyphosate and their interaction to earthworm (*Eisenia fetida*)**  
C.-F. Zhou, Y.-J. Wang, C.-C. Li, R.-J. Sun, Y.-C. Yu, D.-M. Zhou  
*The presence of glyphosate could reduce the toxicity and bioavailability of Cu in the soil because of its strong chelating effects.*

Continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information-Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; Thomson Scientific GeoSciTech; Science Citation Index; SciSearch). Also covered in the abstract and citation database SciVerse Scopus®. Full text available on SciVerse ScienceDirect®.



0269-7491(201309)180;1-H



ELSEVIER

## CONTENTS—Continued from outside back cover

- 78 Micro x-ray absorption spectroscopic analysis of arsenic localization and biotransformation in *Chironomus riparius* Meigen (Diptera: Chironomidae) and *Culex tarsalis* Coquillett (Culicidae)**  
C.L. Mogren, S.M. Webb, W.E. Walton, J.T. Trumble  
*Arsenic accumulation and biotransformation in aquatic insects is variable, but the location and speciation of As provides insight into the detoxification mechanisms of aquatic Diptera.*
- 84 Microbial effects on the release and attenuation of arsenic in the shallow subsurface of a natural geochemical anomaly**  
P. Drahota, L. Falteisek, A. Redlich, J. Rohovec, T. Matoušek, I. Čepička  
*Microbial reduction of Fe(III) and As(V) minerals leads to mobilization of As and induces a mineralogical transition toward realgar formation.*
- 92 Potentially toxic contamination of sediments, water and two animal species in Lake Kalimanci, FYR Macedonia: Relevance to human health**  
P. Vrhovnik, J.P. Arrebola, T. Serafimovski, T. Dolenc, N.R. Šmuc, M. Dolenc, E. Mutch  
*In the Makedonska Kamenica region had been described several potential sources of exposure therefore exists the potential threat to human health.*
- 101 Photochemical trajectory modeling of ozone concentrations in Hong Kong**  
H.R. Cheng, S.M. Saunders, H. Guo, P.K.K. Louie, F. Jiang  
*Isoprene, ethene, toluene, formaldehyde, and m,p-xylene were the key O<sub>3</sub> precursors in Hong Kong.*
- 111 Volatile organic compounds (VOCs) in air from Nisyros Island (Dodecanese Archipelago, Greece): Natural versus anthropogenic sources**  
F. Tassi, F. Capecciacci, L. Giannini, G.E. Vougioukalakis, O. Vaselli  
*Anthropogenic and natural VOCs in air are distinguished on the basis of aromatic, O-substituted, S-substituted and halogenated compounds.*
- 122 Using a toxicokinetics approach to explain the effect of soil pH on cadmium bioavailability to *Folsomia candida***  
M.M. Ardestani, C.A.M. van Gestel  
*Integrating bioaccumulation kinetics with a BLM approach provides novel insights into the bioavailability of cadmium to springtails in soil.*
- 131 Effects of copper on invertebrate–sediment interactions**  
E.R. Hunting, C. Mulder, M.H.S. Kraak, A.M. Breure, W. Admiraal  
*Sub-lethal copper concentrations reduced the bioturbating activities of *A. aquaticus* and invertebrate effects on decomposition, bacterial activity and community metabolic diversity.*
- 136 Matching metal pollution with bioavailability, bioaccumulation and biomarkers response in fish (*Centropomus parallelus*) resident in neotropical estuaries**  
I.C. Souza, I.D. Duarte, N.Q. Pimentel, L.D. Rocha, M. Morozesk, M.M. Bonomo, V.C. Azevedo, C.D.S. Pereira, M.V. Monferrán, C.R.D. Milanez, S.T. Matsumoto, D.A. Wunderlin, M.N. Fernandes  
*Chemometrics allows matching pollution with bioaccumulation of metals and biomarkers responses in the fish *Centropomus parallelus* evidencing differences in estuaries quality.*
- 145 Identification of soil bacteria able to degrade phenanthrene bound to a hydrophobic sorbent *in situ***  
R.K. Regonne, F. Martin, A. Mbawala, M.B. Ngassoum, Y. Jouanneau  
*Bacteria able to degrade PAHs bound to a hydrophobic sorbent were mainly identified as uncultured Rhodocyclaceae and Sphingomonadaceae in polluted soils from tropical and temperate area, respectively.*
- 152 Limited transport of functionalized multi-walled carbon nanotubes in two natural soils**  
D. Kasel, S.A. Bradford, J. Šimůnek, T. Pütz, H. Vereecken, E. Klumpp  
*In undisturbed columns and a lysimeter study, complete retention of functionalized multi-walled carbon nanotubes was found in two soils at environmentally relevant conditions.*
- 159 Organophosphorus esters in the oceans and possible relation with ocean gyres**  
W. Cheng, Z. Xie, J.M. Blais, P. Zhang, M. Li, C. Yang, W. Huang, R. Ding, L. Sun  
*Our work provides a circumpolar investigation on OPEs in the Southern Ocean and we suggest a possibility that ocean currents and gyres may act as important roles in global transport of OPEs.*
- 165 Co-contaminants and factors affecting the sorption behaviour of two sulfonamides in pasture soils**  
P. Srinivasan, A.K. Sarmah, M. Manley-Harris  
*Sorption affinity of SMO and SCP are strongly governed by multitude of factors, and variations in these factors can be significant when manure and fertilisers are added to soil.*
- 173 Influence of the cestode *Ligula intestinalis* and the acanthocephalan *Polymorphus minutus* on levels of heat shock proteins (HSP70) and metallothioneins in their fish and crustacean intermediate hosts**  
S.N. Frank, S. Godehardt, M. Nachev, A. Trubiroha, W. Kloas, B. Sures  
*Parasites are capable of affecting host physiology and therefore modulate biomarker responses.*
- 180 Impact of Zn, Cu, Al and Fe on the partitioning and bioaccessibility of <sup>14</sup>C-phenanthrene in soil**  
I.S. Obuekwe, K.T. Semple  
*High concentrations of Cu and Al (500 mg kg<sup>-1</sup>) can retard the sequestration of phenanthrene in soil.*

CONTENTS—Continued from inside back cover

- 190 Dispersal of the radionuclide caesium-137 (<sup>137</sup>Cs) from point sources in the Barents and Norwegian Seas and its potential contamination of the Arctic marine food chain: Coupling numerical ocean models with geographical fish distribution data**  
H.E. Heldal, F. Vikebø, G.O. Johansen  
*A pulse discharge of <sup>137</sup>Cs from K-159 may cause concentrations in muscle of cod up to 63 and 123 Bq/kg fresh weight in the near-surface and near-bottom layer, respectively.*
- 199 GammaProteobacteria as a potential bioindicator of a multiple contamination by polycyclic aromatic hydrocarbons (PAHs) in agricultural soils**  
M. Niepceon, F. Martin-Laurent, M. Crampon, F. Portet-Koltalo, M. Akpa-Vinceslas, M. Legras, D. Bru, F. Bureau, J. Bodilis  
*The abundance of the GammaProteobacteria class increases significantly in response to a spiking by a polycyclic aromatic hydrocarbon mixture.*
- 206 Interactions of tetracycline with Cd (II), Cu (II) and Pb (II) and their cosorption behavior in soils**  
Y. Zhao, Y. Tan, Y. Guo, X. Gu, X. Wang, Y. Zhang  
*The interaction between tetracycline and metal cations can influence the adsorption behavior of these species in soils, depending on their complexation ability.*
- 214 Oxidation of nonylphenol and octylphenol by manganese dioxide: Kinetics and pathways**  
Z. Lu, J. Gan  
*Naturally occurring MnO<sub>2</sub> may contribute significantly to the attenuation of nonylphenol and octylphenol in soil, water and sediment.*
- 221 Upwind impacts of ammonia from an intensive poultry unit**  
L. Jones, M.S. Nizam, B. Reynolds, S. Bareham, E.R.B. Oxley  
*Ammonia from a poultry unit has upwind impacts, exceeding critical levels 800 m and critical loads 2.8 km upwind, and increasing biomass and tissue N of dune grassland species.*
- 229 Contribution of a submerged membrane bioreactor in the treatment of synthetic effluent contaminated by Bisphenol-A: Mechanism of BPA removal and membrane fouling**  
B. Seyhi, P. Drogui, G. Buelna, A. Azais, M. Heran  
*High BPA removal rates (up to 99%) were obtained in the MBR, with a limit of the toxicity closed to 21.6 g/m<sup>3</sup>/d of BPA.*
- 236 An intermediate complexity dynamic model for predicting accumulation of atmospherically-deposited metals (Ni, Cu, Zn, Cd, Pb) in catchment soils: 1400 to present**  
S. Lofts, E. Tipping, A.J. Lawlor, L. Shotbolt  
*The Intermediate Dynamic Model for Metals predicts the present day pools of Ni, Cu, Zn, Cd and Pb in semi-natural catchment soils of the UK.*
- 246 Health effects of daily airborne particle dose in children: Direct association between personal dose and respiratory health effects**  
G. Buonanno, G.B. Marks, L. Morawska  
*The respiratory health effects of daily airborne particle dose on children through personal monitoring.*
- 251 Bridging the gap between traffic generated health stressors in urban areas: Predicting xylene levels in EU cities**  
C. Vlachokostas, A.V. Michailidou, D. Spyridi, N. Moussiopoulos  
*The multi-stressor, multi-city stepwise regression modelling approach develops reliable statistical associations which capture m,p-xylene and o-xylene trends in EU traffic intensive environments.*
- 259 Historical arsenic contamination of soil due to long-term phosphate fertiliser applications**  
T.N. Hartley, A.J. Macdonald, S.P. McGrath, F.-J. Zhao  
*Long-term phosphate applications before 1948 caused arsenic contamination in the soil of the Park Grass Experiment.*
- 265 Soil contamination by phthalate esters in Chinese intensive vegetable production systems with different modes of use of plastic film**  
J. Wang, Y. Luo, Y. Teng, W. Ma, P. Christie, Z. Li  
*PAE contamination of intensively managed vegetable soils varied widely depending on the mode of use of plastic film in different production systems.*
- 274 Quantitation of tetrabromobisphenol-A from dust sampled on consumer electronics by dispersed liquid-liquid microextraction**  
G. Di Napoli-Davis, J.E. Owens  
*This work describes the analysis of the brominated flame retardant, tetrabromobisphenol-A (TBBPA), from dust sampled on surfaces of consumer electronics.*
- 281 Dioxin-induced acute cardiac mitochondrial oxidative damage and increased activity of ATP-sensitive potassium channels in Wistar rats**  
S.P. Pereira, G.C. Pereira, C.V. Pereira, F.S. Carvalho, M.H. Cordeiro, P.C. Mota, J. Ramalho-Santos, A.J. Moreno, P.J. Oliveira  
*Dioxin alters the regulation of cardiac mitochondrial ATP-sensitive potassium channels and disturbs mitochondrial physiology.*
- 291 Prenatal methylmercury exposure through maternal rice ingestion: Insights from a feasibility pilot in Guizhou Province, China**  
S.E. Rothenberg, X. Yu, Y. Zhang  
*Within a population where rice is a staple food, maternal blood may be an appropriate biomarker for prenatal methylmercury exposure.*
- 299 Effects of a three-year exposure to ambient ozone on biomass allocation in poplar using ethylenediurea**  
Y. Hoshika, F. Pecori, I. Conese, T. Bardelli, E. Marchi, W.J. Manning, O. Badea, E. Paoletti  
*Ozone reduced allocation of biomass to leaves, roots and lateral branches, but did not affect stem length and diameter.*

- 304 Identification of environmental lead sources and pathways in a mining and smelting town: Mount Isa, Australia**  
A.K. Mackay, M.P. Taylor, N.C. Munksgaard, K.A. Hudson-Edwards, L. Burn-Nunes  
*Environmental Pb contamination in the Mount Isa urban area is derived primarily from smelter emissions and local fugitive mining sources and not natural in situ surface mineralization.*
- 312 Filling the gap: Estimating physicochemical properties of the full array of polybrominated diphenyl ethers (PBDEs)**  
C. Yue, L.Y. Li  
*Through careful selection of literature data, structure–property estimation and adjustment, key properties of 209 PBDE congeners are estimated with internal consistency.*
- 324 Review on the effects of toxicants on freshwater ecosystem functions**  
K. Peters, M. Bundschuh, R.B. Schäfer  
*Concentrations assumed as protective in chemical regulation cause adverse effects in three fundamental ecosystem functions.*
- 330 Guidelines for biomonitoring persistent organic pollutants (POPs), using lichens and aquatic mosses – A review**  
S. Augusto, C. Máguas, C. Branquinho  
*This review fulfils the lack of knowledge regarding the use of lichens and aquatic mosses as biomonitors of POPs, providing a set of guidelines to be followed.*
- 339 Ecological risk of estrogenic endocrine disrupting chemicals in sewage plant effluent and reclaimed water**  
Y. Sun, H. Huang, Y. Sun, C. Wang, X.-L. Shi, H.-Y. Hu, T. Kameya, K. Fujie  
*The results suggested that 17 $\alpha$ -ethynylestradiol (EE2), estrone (E1) and estradiol (E2) should be the priority EDCs to control in municipal sewage plants.*
- 345 Toxicities of oils, dispersants and dispersed oils to algae and aquatic plants: Review and database value to resource sustainability**  
M. Lewis, R. Pryor  
*The phytotoxicity database does not support toxicity predictions for most oils and dispersants.*
- 368 Biogas final digestive byproduct applied to croplands as fertilizer contains high levels of steroid hormones**  
C. Rodriguez-Navas, E. Björklund, B. Halling-Sørensen, M. Hansen  
*Eight steroid hormones were found in biogas digestate byproduct in the ng g<sup>-1</sup> dm levels. Anaerobic digestion processes do not completely remove steroid hormones from organic waste residues.*
- 372 What is required for the validation of in vitro assays for predicting contaminant relative bioavailability? Considerations and criteria**  
A.L. Juhasz, N.T. Basta, E. Smith  
*Defined criteria for establishing in vivo–in vitro performance validation are required in order to ensure robust, defensible predictive models for human health exposure assessment.*