

TU  
E58



ELSEVIER

Volume 182

November 2013

ISSN 0269-7491

# ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

**W.J. Manning**


ASSOCIATE EDITORS

**J. Gan**

**B. Nowack**

**E. Paoletti**

**C. Wiegand**



## CONTENTS

Volume 182 November 2013

- 1 Occurrence and distribution of pharmaceutically active and endocrine disrupting compounds in Singapore's marine environment: Influence of hydrodynamics and physical-chemical properties**  
S. Bayen, H. Zhang, M.M. Desai, S.K. Ooi, B.C. Kelly  
*A field study demonstrates the influence of hydrodynamic residence time and physical-chemical properties on exposure risks of PhACs and EDCs in coastal marine ecosystems.*
- 9 Have the short-term mortality effects of particulate matter air pollution changed in Australia over the period 1993–2007?**  
S. Roberts  
*We show that the mortality effect of  $PM_{10}$  has declined in Australia over the period 1993–2007, possibly due to a reduction in the toxicity of  $PM_{10}$ .*
- 15 Concentrations and geographic distribution of selected organic pollutants in Scottish surface soils**  
S.M. Rhind, C.E. Kyle, C. Kerr, M. Osprey, Z.L. Zhang, E.I. Duff, A. Lilly, A. Nolan, G. Hudson, W. Towers, J. Bell, M. Coull, C. McKenzie  
*Soil concentrations of anthropogenic persistent organic pollutants are not clearly related to soil type or pH, vegetation, altitude, or distance from pollutant sources.*
- 28 Influence of urbanization and industrialization on metal enrichment of sediment cores from Shantou Bay, South China**  
Y. Qiao, Y. Yang, J. Zhao, R. Tao, R. Xu  
*Shantou Bay was polluted by Cd, Cu, Pb and Zn, and showed an increase trend along time. River input, metropolitan and port facilities were identified as their sources based on multi-analysis.*
- 37 The influence of dissolved phosphorus molecular form on recalcitrance and bioavailability**  
B. Li, M.T. Brett  
*SRP is a poor indicator of the bioavailability of many of P containing compounds and much of what is classified as SRP in nature could be associated with humic-metal complexes with low bioavailability.*
- 45 Influence of relative trophic position and carbon source on selenium bioaccumulation in turtles from a coal fly-ash spill site**  
J.U. Van Dyke, W.A. Hopkins, B.P. Jackson  
*Stable isotope differences indicate that claw selenium concentrations differ among relative carbon sources, and not among relative trophic positions, in an assemblage of aquatic turtles.*
- 53 Enhancing the biodegradation of oil in sandy sediments with choline: A naturally methylated nitrogen compound**  
B. Mortazavi, A. Horel, J.S. Anders, A. Mirjafari, M.J. Beazley, P.A. Sobczyk  
*Choline, a naturally occurring methylated nitrogen-containing compound, accelerated hydrocarbon degradation in sandy sediments by an extant microbial community.*
- 63 Placental IGF-1 and IGFBP-3 expression correlate with umbilical cord blood PAH and PBDE levels from prenatal exposure to electronic waste**  
X. Xu, T.A. Yekeen, Q. Xiao, Y. Wang, F. Lu, X. Huo  
*The correlation between POPs and placental IGF-1 and IGFBP-3 assessed provides more information on the human health risk associated with electronic waste.*
- 70 Critical comparison of intravenous injection of  $TiO_2$  nanoparticles with waterborne and dietary exposures concludes minimal environmentally-relevant toxicity in juvenile rainbow trout *Oncorhynchus mykiss***  
D. Boyle, G.A. Al-Bairuty, T.B. Henry, R.D. Handy  
*Critical evaluation of directly comparable investigations of  $TiO_2$ -NP toxicity by waterborne, dietary, and intravenous injection exposures conclude minimal toxicity in juvenile rainbow trout.*

Continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/ Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information-Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; Thomson Scientific GeoSciTech; Science Citation Index; SciSearch). Also covered in the abstract and citation database Scopus®. Full text available on ScienceDirect®.



0269-7491(201311)182;1-K



ELSEVIER

## CONTENTS—Continued from outside back cover

- 80 A multi-isotope approach for estimating industrial contributions to atmospheric nitrogen deposition in the Athabasca oil sands region in Alberta, Canada**  
B.C. Proemse, B. Mayer, M.E. Fenn, C.S. Ross  
*Distinct  $\delta^{18}\text{O}$ ,  $\Delta^{17}\text{O}$ , and  $\delta^{15}\text{N}$  values were used to estimate industrially derived N contributions to atmospheric nitrate and ammonium deposition in the Athabasca oil sands region.*
- 92 In-situ assessment of metal contamination via portable X-ray fluorescence spectroscopy: Zlatna, Romania**  
D.C. Weindorf, L. Paulette, T. Man  
*Metal contamination in Zlatna, Romania exceeds regulatory limits; portable X-ray fluorescence spectrometry and kriging interpolation were used to model contaminants in-situ.*
- 101 Long-term trend of haze pollution and impact of particulate matter in the Yangtze River Delta, China**  
Z. Cheng, S. Wang, J. Jiang, Q. Fu, C. Chen, B. Xu, J. Yu, X. Fu, J. Hao  
*The long-term variation of haze pollution in the YRD and its cause is investigated and the quantitative contribution of particulate matter to haze pollution is estimated.*
- 111 Assessing mixed trace elements in groundwater and their health risk of residents living in the Mekong River basin of Cambodia**  
K. Phan, S. Phan, L. Huoy, B. Suy, M.H. Wong, J.H. Hashim, M.S. Mohamed Yasin, S.M. Aljunid, S. Sthiannopkao, K.-W. Kim  
*Risk assessment indicated that 98.7% of residents in Kandal and 12.4% of Kratie study areas were at risk of non-carcinogenic effects of multiple elements in groundwater.*
- 120 Polybrominated diphenyl ethers affect the reproduction and development, and alter the sex ratio of zebrafish (*Danio rerio*)**  
X.B. Han, K.W.Y. Yuen, R.S.S. Wu  
*PBDE reduces fertilization and hatching successes, causes malformation and leads to a male biased F1 generation in fish.*
- 127 Spatial and vertical distribution of mercury in upland forest soils across the northeastern United States**  
J.B. Richardson, A.J. Friedland, T.R. Engerbretson, J.M. Kaste, B.P. Jackson  
*Forest floor Hg was correlated with soil organic matter, pH, latitude, and mean annual precipitation. Mineral soil Hg was greatest in Bs horizons of Spodosols.*
- 135 A method to analyze "source-sink" structure of non-point source pollution based on remote sensing technology**  
M. Jiang, H. Chen, Q. Chen  
*"Source-Sink" Structure of non-point source nitrogen and phosphorus pollution in Jiulongjiang estuary in China was worked out by the Grid Landscape Contrast Index.*
- 141 Behavior of Ag nanoparticles in soil: Effects of particle surface coating, aging and sewage sludge amendment**  
A.R. Whitley, C. Levard, E. Oostveen, P.M. Bertsch, C.J. Matocha, F.v.d. Kammer, J.M. Unrine  
*Behavior of manufactured silver nanoparticles in soil depends on surface coating, contact with sewage sludge, and aging.*
- 150 Uptake and accumulation of four PPCP/EDCs in two leafy vegetables**  
L.K. Dodgen, J. Li, D. Parker, J.J. Gan  
*Four commonly occurring PPCP/EDCs were taken up into roots of lettuce and collards, but translocation into edible leaves was limited and nearly all residue was non-extractable.*
- 157 Spatio-temporal variation of biogenic volatile organic compounds emissions in China**  
L.Y. Li, Y. Chen, S.D. Xie  
*An emission inventory of BVOCs in China was established based on the most detailed and latest vegetation investigations, and high-resolution meteorological data.*
- 169 Adsorption of carbamazepine by carbon nanotubes: Effects of DOM introduction and competition with phenanthrene and bisphenol A**  
I. Lerman, Y. Chen, B. Xing, B. Chefetz  
*In multi-component system including the main adsorbate and competitor, DOM exhibited significant effect on adsorption of contaminants by carbon nanotubes.*
- 177 Abiotic degradation of antibiotic ionophores**  
P. Bohn, S.A. Bak, E. Björklund, K.A. Krogh, M. Hansen  
*Antibiotic ionophores were found to undergo either hydrolysis in acidic environments (monensin, salinomycin, and narasin) or photolysis (lasalocid).*
- 184 Levels of prenatal mercury exposure and their relationships to neonatal anthropometry in Wujiang City, China**  
B.-Q. Guo, S.-Z. Cai, J.-L. Guo, J. Xu, W. Wu, H. Li, X. Zhou, D.-S. Kim, C.-H. Yan, H.-D. Lü  
*The levels of prenatal Hg exposure were low and no significant correlation was found between low-level Hg exposure and neonatal anthropometry in Wujiang City, China.*
- 190 Biogeochemistry and community ecology in a spring-fed urban river following a major earthquake**  
N.S. Wells, T.J. Clough, L.M. Condron, W.T. Baisden, J.S. Harding, Y. Dong, G.D. Lewis, G. Lear  
*Linking stream community ecology with biogeochemical function, we provide an in-depth quantification of urban stream recovery following a catastrophic earthquake.*

**CONTENTS—Continued from inside back cover**

- 201 *In situ* application of activated carbon and biochar to PCB-contaminated soil and the effects of mixing regime**  
M.J. Denyes, A. Rutter, B.A. Zeeb  
*In situ AC and biochar soil amendments perform equally well at reducing PCB uptake, however, laboratory-based mixing methods may exaggerate the sorptive capacities of both amendments.*
- 209 Ultrastructural changes and Heat Shock Proteins 70 induced by atmospheric pollution are similar to the effects observed under *in vitro* heavy metals stress in *Conocephalum conicum* (Marchantiales – Bryophyta)**  
A. Basile, S. Sorbo, B. Conte, M. Cardi, S. Esposito  
*Pollution induces ultrastructural changes and Hsp70 induction in C. conicum, that can be mimicked in vitro. This liverwort could be used as bioindicator.*
- 217 Mineralogy affects geoavailability, bioaccessibility and bioavailability of zinc**  
R.M. Molina, L.A. Schaidler, T.C. Donaghey, J.P. Shine, J.D. Brain  
*Zinc mineralogy influences in vitro bioaccessibility and in vivo bioavailability and in vitro extraction tests can be used to predict Zn bioavailability from particles.*
- 225 Effects of local and spatial conditions on the quality of harvested rainwater in the Mekong Delta, Vietnam**  
G.-J. Wilbers, Z. Sebesvari, A. Rechenburg, F.G. Renaud  
*Concentrations of lead and total coliforms in household-harvested rainwater in the Mekong Delta exceed drinking water guidelines in 17% and 92% of the samples, respectively.*
- 233 Impact of carbonaceous materials in soil on the transport of soil-bound PAHs during rainfall-runoff events**  
X. Luo, Y. Zheng, B. Wu, Z. Lin, F. Han, W. Zhang, X. Wang  
*Anthropogenic carbonaceous materials in soil, especially black carbon, largely control the transport of soil-bound PAHs during rainfall-runoff events.*
- 242 Photosynthetic response of early and late leaves of white birch (*Betula platyphylla* var. *japonica*) grown under free-air ozone exposure**  
Y. Hoshika, M. Watanabe, N. Inada, Q. Mao, T. Koike  
*Early leaves have higher resistance to ozone stress than late leaves in heterophyllous white birch.*
- 248 Complexation with dissolved organic matter and mobility control of heavy metals in the rhizosphere of hyperaccumulator *Sedum alfredii***  
T. Li, Q. Tao, C. Liang, M.J.I. Shohag, X. Yang, D.L. Sparks  
*DOM in the rhizosphere of hyperaccumulator S. alfredii could significantly increase metal mobility through the formation of soluble DOM-metal complexes.*
- 256 Mercury concentrations in human placenta, umbilical cord, cord blood and amniotic fluid and their relations with body parameters of newborns**  
I. Kozikowska, Ł.J. Binkowski, K. Szczepańska, H. Sławska, K. Miszczuk, M. Śliwińska, T. Łaciak, R. Stawarz  
*Maternal age was not an influential factor of mercury concentrations in studied samples. 75% of cord blood samples exceeded the Hg threshold concentration.*
- 263 Comparison of earthworm responses to petroleum hydrocarbon exposure in aged field contaminated soil using traditional ecotoxicity endpoints and <sup>1</sup>H NMR-based metabolomics**  
M. Whitfield Åslund, G.L. Stephenson, A.J. Simpson, M.J. Simpson  
*Earthworm metabolomic responses measured after 2 and 14 days are compared to traditional earthworm ecotoxicity endpoints (survival and reproduction) in petroleum hydrocarbon contaminated soil.*
- 269 The effects of surfactants and solution chemistry on the transport of multiwalled carbon nanotubes in quartz sand-packed columns**  
Y. Lu, X. Xu, K. Yang, D. Lin  
*The MWCNT transport through porous media varied with surfactant property and solution chemistry.*
- 278 Associations of particulate air pollution and daily mortality in 16 Chinese cities: An improved effect estimate after accounting for the indoor exposure to particles of outdoor origin**  
R. Chen, B. Zhou, H. Kan, B. Zhao  
*Accounting for the indoor PM<sub>10</sub> concentrations of outdoor origin in the exposure assessment could improve the effect estimates of ambient PM<sub>10</sub>.*
- 283 Atmospheric polychlorinated biphenyls in Indian cities: Levels, emission sources and toxicity equivalents**  
P. Chakraborty, G. Zhang, S. Eckhardt, J. Li, K. Breivik, P.K.S. Lam, S. Tanabe, K.C. Jones  
*Measurement of atmospheric Polychlorinated biphenyls in seven major Indian cities.*
- 291 Influence of socio-demographic and diet determinants on the levels of mercury in preschool children from a Mediterranean island**  
M. Garí, J.O. Grimalt, M. Torrent, J. Sunyer  
*Oily fish and shellfish consumption, parity, maternal smoking and occupational status are the main determinants of mercury in four year-old children from Menorca Island (Mediterranean Sea).*
- 299 Wet deposition of brominated flame retardants to the Great Lakes basin – Status and trends**  
M. Robson, L. Melymuk, L. Bradley, B. Treen, S. Backus  
*Spatial and temporal trends in the wet deposition of brominated flame retardants reveal a decline in BDE 209 concentrations and the importance of episodic high concentration events.*

- 307 Localized enrichment of polycyclic aromatic hydrocarbons in soil, spruce needles, and lake sediments linked to *in-situ* bitumen extraction near Cold Lake, Alberta**  
J.B. Korosi, G. Irvine, E.K. Skierszkan, J.R. Doyle, L.E. Kimpe, J. Janvier, J.M. Blais  
*PAHs in sediments and soils were generally low in areas adjacent to in-situ bitumen extraction rigs in the Cold Lake Alberta oil sands, but evidence of localized contamination at some sites was evident.*
- 316 Familial differences in the effects of mercury on reproduction in zebra finches**  
C.W. Varian-Ramos, J.P. Swaddle, D.A. Cristol  
*Genetic variation in response to contaminants can lead to adaptation on long-term contaminated sites, with implications for risk assessment and conservation of impacted populations.*
- 324 Source contribution analysis of surface particulate polycyclic aromatic hydrocarbon concentrations in northeastern Asia by source-receptor relationships**  
Y. Inomata, M. Kajino, K. Sato, T. Ohara, J.-i. Kurokawa, H. Ueda, N. Tang, K. Hayakawa, T. Ohizumi, H. Akimoto  
*Transboundary transport of PAHs in northeast Asia was investigated by source-receptor analysis.*
- 335 Accumulation of wet-deposited radiocaesium and radiostrontium by spring oilseed rape (*Brassica napus* L.) and spring wheat (*Triticum aestivum* L.)**  
Stefan.B. Bengtsson, J. Eriksson, A.I. Gärdenäs, M. Vinichuk, K. Rosén  
*Intercepted radionuclides can be transferred into the food chain for humans.*
- 343 Ozone distribution in remote ecologically vulnerable terrain of the southern Sierra Nevada, CA**  
J. Panek, D. Saah, A. Esperanza, A. Bytnerowicz, W. Fraczek, R. Cisneros  
*A passive ozone sampler network in combination with spatial analysis techniques was used to characterize landscape-scale ozone patterns and dynamics, identifying regions of consistently high and low ozone exposure for forest management prioritization.*
- 357 Sediment pore water distribution coefficients of PCB congeners in enriched black carbon sediment**  
A. Martinez, C. O'Sullivan, D. Reible, K.C. Hornbuckle  
*The organic carbon fraction times the octanol-water partition coefficient yielded the best prediction model for the sediment pore water distribution coefficient of PCBs.*
- 364 Space and time resolved monitoring of airborne particulate matter in proximity of a traffic roundabout in Sweden**  
K.E. Wilkinson, J. Lundkvist, J. Netrval, M. Eriksson, G.A. Seisenbaeva, V.G. Kessler  
*Automated SEM-EDS analysis of captured roadside PM at a traffic roundabout in Sweden displaying the time- and space-resolved chemical differences of the captured particles.*
- 371 Adverse effects induced by ecgonine methyl ester to the zebra mussel: A comparison with the benzoyllecgonine**  
M. Parolini, A. Binelli  
*Environmentally relevant ecgonine methyl ester concentrations induced adverse effects to zebra mussels.*
- 379 Estimating the toxicity of ambient fine aerosols using freshwater rotifer *Brachionus calyciflorus* (Rotifera: Monogononta)**  
V. Verma, R. Rico-Martinez, N. Kotra, C. Rennolds, J. Liu, T.W. Snell, R.J. Weber  
*The toxicity of ambient fine aerosols as estimated using freshwater rotifers is largely contributed by their hydrophobic components.*
- 385 Study of molecularly imprinted solid-phase extraction of gonyautoxins 2,3 in the cultured dinoflagellate *Alexandrium tamarense* by high-performance liquid chromatography with fluorescence detection**  
Z.-R. Lian, J.-T. Wang  
*MIPMs for GTX<sub>2,3</sub> were prepared and applied as special SPE sorbents. The MISPE process was valid for the isolation and clean-up of GTX<sub>2,3</sub> from *A. tamarense* extract.*
- 392 Spatio-temporal variations of black carbon concentrations in the Megacity Beijing**  
N. Schleicher, S. Norra, M. Fricker, U. Kaminski, Y. Chen, F. Chai, S. Wang, Y. Yu, K. Cen  
*Black carbon was investigated for a two-year period in the megacity Beijing to gain detailed knowledge about the seasonal, temporal and spatial patterns of BC particles and their sources.*
- 402 Antibiotics in riverine runoff of the Pearl River Delta and Pearl River Estuary, China: Concentrations, mass loading and ecological risks**  
W. Xu, W. Yan, X. Li, Y. Zou, X. Chen, W. Huang, L. Miao, R. Zhang, G. Zhang, S. Zou  
*Antibiotics were ubiquitous in the river and coastal water in the Pearl River Delta and posed various ecological risks to the relevant aquatic organisms.*
- 408 Distributions, sources and pollution status of 17 trace metal/metalloids in the street dust of a heavily industrialized city of central China**  
Z. Li, X. Feng, G. Li, X. Bi, J. Zhu, H. Qin, Z. Dai, J. Liu, Q. Li, G. Sun  
*Pb/Zn smelting and hard alloy processing operations have caused seriously contamination of trace metal/metalloids in the street dust.*
- 417 Improved retrieval of PM<sub>2.5</sub> from satellite data products using non-linear methods**  
M. Sorek-Hamer, A.W. Strawa, R.B. Chatfield, R. Esswein, A. Cohen, D.M. Broday  
*Improved PM<sub>2.5</sub> retrievals from satellite observations can fill in missing surface measurements in areas with sparse ground monitoring and be used for evaluating air quality models and as exposure metrics.*
- 424 Polychlorinated biphenyls in surface soil in urban and background areas of Mongolia**  
E.A. Mamontova, A.A. Mamontov, E.N. Tarasova, M.I. Kuzmin, D. Ganchimeg, M.Yu. Khomutova, O. Gombosuren, E. Ganjuurjav  
*Polychlorinated biphenyls were measured in soils throughout Mongolia.*
- 430 Atmospheric organic nitrogen deposition: Analysis of nationwide data and a case study in Northeast China**  
C.M. Jiang, W.T. Yu, Q. Ma, Y.G. Xu, H. Zou, S.C. Zhang, W.P. Sheng  
*Synthesis of DON deposition across China implied regional importance of anthropogenic sources, and an observation in Northeast China suggested the ecological significances of the DON flux should be considered.*

- 437 Modelling relationships between lichen bioindicators, air quality and climate on a national scale: Results from the UK OPAL air survey**  
L. Seed, P. Wolseley, L. Gosling, L. Davies, S.A. Power  
*Data on selected lichen taxa collected by members of the public in England is used to show the relationship of indicator taxa and pollution indices to air pollution and climate data.*
- 448 Can citizen science produce good science? Testing the OPAL Air Survey methodology, using lichens as indicators of nitrogenous pollution**  
D.J. Tregidgo, S.E. West, M.R. Ashmore  
*A simplified lichen biomonitoring method used for citizen science can detect the impact of nitrogenous air pollution from local roads.*
- 452 Biphasic dose responses in biology, toxicology and medicine: Accounting for their generalizability and quantitative features**  
E.J. Calabrese  
*This paper provides a biologically based explanation for the generalizability/quantitative features of the hormetic dose response, representing a fundamental contribution to the field.*
- 461 Sources, factors, mechanisms and possible solutions to pollutants in marine ecosystems**  
K.M.G. Mostofa, C.-Q. Liu, D. Vione, K. Gao, H. Ogawa  
*Review of sources, factors, mechanisms and possible remedial measures of key pollutants (contaminants, toxins, ship breaking, overfishing) in marine ecosystems.*
- 479 Pyrethroid effects on freshwater invertebrates: A meta-analysis of pulse exposures**  
J.J. Rasmussen, P. Wiberg-Larsen, E.A. Kristensen, N. Cedergreen, N. Friberg  
*This review shows that pulse exposure of pyrethroids have long-term effects on macroinvertebrates at lower concentrations than 1/100 of the 48 h LC50 for D. magna.*
- 486 Photobleaching of lignin derived compounds from pulp mill effluents upon irradiation: The key role of receiving waters**  
M. Otero, I. Guilherme, E.B.H. Santos  
*Under irradiation, DOM from kraft pulp mill effluents progressively resembles that from not-polluted sites, which may not occur if effluents are discharged in saline waters.*
- 490 Anthropogenic contaminants in Indo-Pacific humpback and Australian snubfin dolphins from the central and southern Great Barrier Reef**  
D. Cagnazzi, M.C. Fossi, G.J. Parra, P.L. Harrison, S. Maltese, D. Coppola, A. Soccodato, M. Bent, L. Marsili  
*Potentially hazardous levels of some coastal contaminants were found in two species of dolphins inhabiting the Great Barrier Reef Marine Park coastal region.*
- 495 Microplastic pollution in deep-sea sediments**  
L. Van Cauwenberghe, A. Vanreusel, J. Mees, C.R. Janssen  
*Here, we demonstrate that microplastics have invaded the marine environment to an extent that they appear to even be present in the remote deep sea.*

#### Letter to the Editor

- 500 Comments on "Characterising metal build-up on urban road surfaces" by Egodawatta et al. (2013). Environmental Pollution 176, 87–91**  
J. Zhang, P. Krebs

#### Reply to Letter to the Editor

- 503 Reply to comment on "Characterising metal build-up on urban road surfaces" by Egodawatta et al. (2013). Environmental Pollution, 176, 87–91**  
P. Egodawatta, A.M. Ziyath, A. Goonetilleke