


7M

E58

Volume 189

June 2014


ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning



CONTENTS

Volume 189 June 2014

- 1 Response of detoxification gene mRNA expression and selection of molecular biomarkers in the clam *Ruditapes philippinarum* exposed to benzo[a]pyrene**
D. Liu, L. Pan, Y. Cai, Z. Li, J. Miao
mRNA expressions of AhR, GST-pi and Mn-SOD in gills and GST-pi in digestive glands of the clams are useful molecular biomarkers of B[a]P exposure.
- 9 Effects of elevated ozone, carbon dioxide, and the combination of both on the grain quality of Chinese hybrid rice**
Y. Wang, Q. Song, M. Frei, Z. Shao, L. Yang
Exposure of Chinese hybrid rice to elevated ozone and CO₂ during growth causes fewer changes in grain quality than ozone exposure alone.
- 18 Chemical cocktails in aquatic systems: Pesticide effects on the response and recovery of >20 animal taxa**
J. Hua, R. Relyea
Insecticides applied individually and in a mixture have complex direct and indirect consequences on aquatic system response and recovery.
- 27 Bioaccumulation of perfluoroalkyl compounds in midge (*Chironomus riparius*) larvae exposed to sediment**
D. Bertin, B.J.D. Ferrari, P. Labadie, A. Sapin, J. Garric, H. Budzinski, M. Houde, M. Babut
Chironomus riparius mainly bioaccumulates long-chain PFASs via trophic and/or tegumentary routes during the fourth instar larvae growth phase.
- 35 Reactivity and fate of secondary alkane sulfonates (SAS) in marine sediments**
R.M. Baena-Nogueras, P. Rojas-Ojeda, J.L. Sanz, E. González-Mazo, P.A. Lara-Martín
Anaerobic degradation of SAS in marine sediments and the metabolites involved in this process are reported for the first time.
- 43 Comparative use of lichens, mosses and tree bark to evaluate nitrogen deposition in Germany**
S.H. Boltersdorf, R. Pesch, W. Werner
First nationwide comparison of lichens, mosses and tree bark to assess the N deposition in Germany by analysing N content and $\delta^{15}N$ values.
- 54 Metal and nutrient dynamics in decomposing tree litter on a metal contaminated site**
L. Van Nevel, J. Mertens, A. Demey, A. De Schrijver, S. De Neve, F.M.G. Tack, K. Verheyen
Litter decomposition rates, as well as enrichment and release dynamics of metals and nutrients in decomposing litter were divergent under the different tree species.
- 63 On the link between biomagnetic monitoring and leaf-deposited dust load of urban trees: Relationships and spatial variability of different particle size fractions**
J. Hofman, K. Wuyts, S. Van Wittenberghe, M. Brackx, R. Samson
The relation between leaf-deposited particle mass and SIRM illustrates the relevancy of biomagnetic monitoring for the quantification of urban particulates.
- 73 Limited effect of urban tree vegetation on NO₂ and O₃ concentrations near a traffic route**
M. Grundström, H. Pleijel
Urban vegetation had limited effect on local air pollution.
- 77 Cumulative risk assessment for plasticizer-contaminated food using the hazard index approach**
J.W. Chang, B.R. Yan, M.H. Chang, S.H. Tseng, Y.M. Kao, J.C. Chen, C.C. Lee
The health of young Taiwanese may be adversely affected by overexposure of plasticizer-contaminated food.
- 85 Haze in China: Current and future challenges**
M. Li, L. Zhang
This paper analyzed the current condition, the cause of heavy haze and measures to tackle air pollution in China.

Continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/ Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information-Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; Thomson Scientific GeoSciTech; Science Citation Index; SciSearch). Also covered in the abstract and citation database Scopus®. Full text available on ScienceDirect®.



0269-7491(201406)189:1-A



ELSEVIER

CONTENTS—Continued from outside back cover

- 87 Does water chemistry affect the dietary uptake and toxicity of silver nanoparticles by the freshwater snail *Lymnaea stagnalis*?**
A.L.-S. Oliver, M.-N. Croteau, T.L. Stoiber, M. Tejamaya, I. Römer, J.R. Lead, S.N. Luoma
L. Stagnalis assimilated PVP-AgNPs efficiently from food and water chemistry had no influence on their uptake and toxicity.
- 92 Assessing biochar's ability to reduce bioavailability of aminocyclopyrachlor in soils**
J.L. Rittenhouse, P.J. Rice, K.A. Spokas, W.C. Koskinen
This was the first study to assess the use of biochar as a remediation tool for reducing bioavailable aminocyclopyrachlor in the liquid phase soil systems.
- 98 Metals in European roadside soils and soil solution – A review**
M. Werkenthin, B. Kluge, G. Wessolek
Summarised data showed typical distance related metal patterns of European roadside soils; solute concentrations are mostly independent from soil matrix concentrations.
- 111 Highlighting the threat from current and near-future ozone pollution to clover in pasture**
D.K.L. Hewitt, G. Mills, F. Hayes, S. Wilkinson, W. Davies
Ozone effects on the growth and functioning of clover cultivars.
- 118 Intake estimates of phthalate esters for South Delhi population based on exposure media assessment**
M.T. Das, P. Ghosh, I.S. Thakur
DEHP was the most abundant congener and its daily intake reached upto $70 \mu\text{g kg}^{-1} \text{d}^{-1}$. Intake doses of DnBP and DEHP reached levels near or above the exposure limits.
- 126 Do predictions from Species Sensitivity Distributions match with field data?**
S. Smetanová, L. Bláha, M. Liess, R.B. Schäfer, M.A. Beketov
Comparison of the SSD-based prediction with the field data evaluated with the $\text{SPEAR}_{\text{pesticides}}$ index shows that SSD threshold msPAF of 0.05 severely underestimates the effects observed in the field.
- 134 Atrazine contamination at the watershed scale and environmental factors affecting sampling rates of the polar organic chemical integrative sampler (POCIS)**
R.L. Dalton, F.R. Pick, C. Boutin, A. Saleem
POCIS effectively measured atrazine across a watershed, while field calibration indicated that sampling rates were affected by temperature and not other environmental factors.
- 143 Bacterial metal resistance genes and metal bioavailability in contaminated sediments**
S. Roosa, R. Wattiez, E. Prygiel, L. Lesven, G. Billon, D.C. Gillan
*Capsule Bacterial *czcA* is a potential biomarker of Cd, Zn and Co bioavailability in aquatic sediments as shown by quantitative PCR and sequential metal extraction.*
- 152 Combining measurements and modelling to quantify the contribution of atmospheric fallout, local industry and road traffic to PAH stocks in contrasting catchments**
D. Gateuille, O. Evrard, I. Lefevre, E. Moreau-Guigon, F. Alliot, M. Chevreuil, J.-M. Mouchel
Source contributions to PAH soil contamination were investigated with distance-based models.
- 161 High tolerance of subalpine grassland to long-term ozone exposure is independent of N input and climatic drivers**
M. Volk, V. Wolff, S. Bassin, C. Ammann, J. Fuhrer
Subalpine grassland is tolerant to increasing ozone exposure, independent of N input and climatic drivers, while N deposition rates at the critical load strongly increase productivity.
- 169 Characterization of estrogen receptor α activities in polychlorinated biphenyls by *in vitro* dual-luciferase reporter gene assay**
Q. Zhang, M. Lu, C. Wang, J. Du, P. Zhou, M. Zhao
*The agonist and antagonist activities of 20 PCBs were investigated and compared in *in vitro* models.*
- 176 Repeated phytoextraction of four metal-contaminated soils using the cadmium/zinc hyperaccumulator *Sedum plumbizincicola***
Z. Li, L. Wu, P. Hu, Y. Luo, H. Zhang, P. Christie
Acid soil has high total metal phytoremediation efficiency while a strategy based on stripping of the bioavailable contaminant might be feasible for alkaline soil phytoremediation
- 184 Pyruvate carboxylase as a sensitive protein biomarker for exogenous steroid chemicals**
X. Liang, C.-J. Martyniuk, G. Cheng, J. Zha, Z. Wang
Our results suggest PC may be a sensitive biomarker of exposure to exogenous steroid chemicals in the liver of fish.
- 194 Transboundary smoke haze pollution in Malaysia: Inpatient health impacts and economic valuation**
J. Othman, M. Sahani, M. Mahmud, M.K. Sheikh Ahmad
Inpatient rates soared by 31% while economic loss valued at USD91,000 annually.

CONTENTS—Continued from inside back cover

- 202 Effects of the geophagous earthworm *Metaphire guillelmi* on sorption, mineralization, and bound-residue formation of 4-nonylphenol in an agricultural soil**
J. Shan, Y. Wang, L. Wang, X. Yan, R. Ji
Earthworms significantly changed the fate of 4-NP, highlighting that effects of earthworm should be considered when evaluating the behavior and risk of 4-NP in soil.
- 208 Metal composition of fine particulate air pollution and acute changes in cardiorespiratory physiology**
S. Cakmak, R. Dales, L.M. Kauri, M. Mahmud, K. Van Ryswyk, J. Vanos, L. Liu, P. Kumarathasan, E. Thomson, R. Vincent, S. Weichenthal
Several PM_{2.5} metals were associated with acute changes in cardiovascular or respiratory physiology. Given the evidence of source specificity, our study provides novel information.
- 215 Relating metal bioavailability to risk assessment for aquatic species: Daliao River watershed, China**
S. Han, Y. Zhang, S. Masunaga, S. Zhou, W. Naito
The highest bioavailability values and the highest risks of metals were found in the estuary of the Daliao River.
- 223 Lead and other elements in house dust of Japanese residences – Source of lead and health risks due to metal exposure**
J. Yoshinaga, K. Yamasaki, A. Yonemura, Y. Ishibashi, T. Kaido, K. Mizuno, M. Takagi, A. Tanaka
Major source of Pb in house dust of Japanese residences was not track-in soil but unknown materials that contain Pb and Sb and/or Sn.
- 229 The impact of gypsum mine water: A case study on morphology and DNA integrity in the freshwater invertebrate, *Gammarus balcanicus***
I. Ternjej, Z. Mihaljević, M. Ivković, A. Previšić, I. Stanković, K. Maldini, D. Želježić, N. Kopjar
Gypsum mine wastewaters have genotoxic potential and affect the gammarid exoskeleton morphology and biochemistry associated with a high strontium uptake.
- 239 Magnetic susceptibility of road deposited sediments at a national scale – Relation to population size and urban pollution**
D. Jordanova, N. Jordanova, P. Petrov
Magnetic susceptibility of road dusts on a national scale increases proportionally to the population size and mean NO₂ concentrations due to the effect of traffic related pollution.
- Letter to the Editor**
- 252 Combining the effects of conventional risk factors and environmental triggering factors while studying seasonality in acute myocardial infarction**
M. Al Mamun, N. Rumana, Y. Kita, T.C. Turin
Detailed research initiatives are required to get a clear scenario on complete characterization of environmental triggers in relation to conventional cardiovascular risk factors.