


174
E 58

Volume 193

October 2014


ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning



CONTENTS

Volume 193 October 2014

- 1 Gene expression in snapbeans exposed to ozone and protected by ethylenediurea**
E. Paoletti, A. Castagna, L. Ederli, S. Pasqualini, A. Ranieri, W.J. Manning
Transcripts of major antioxidants in a model plant exposed to ozone and protected by EDU were measured for the first time and suggested that EDU may halt the O₃-induced ROS generation.
- 6 Characterization of polychlorinated dibenzo-*p*-dioxins and dibenzofurans, dioxin-like polychlorinated biphenyls, and polychlorinated naphthalenes in the environment surrounding secondary copper and aluminum metallurgical facilities in China**
J. Hu, M. Zheng, W. Liu, Z. Nie, C. Li, G. Liu, K. Xiao
This study suggested that atmospheric PCDD/F, PCB, and PCN contaminations near secondary non-ferrous metal smelters were affected by emissions from the investigated plants.
- 13 Contrasting effects of water salinity and ozone concentration on two cultivars of durum wheat (*Triticum durum* Desf.) in Mediterranean conditions**
G. Gerosa, R. Marzuoli, A. Finco, R. Monga, I. Fusaro, F. Faoro
Durum wheat response to ozone exposure and irrigation water salinity differs in two cultivars known for their different sensitivity to the onset of ozone-like visible leaf injuries.
- 22 The effect of dibenzo-*p*-dioxin- and dibenzofuran-contaminated soil on the earthworm *Eisenia andrei***
R. Roubalová, J. Dvořák, P. Procházková, D. Elhottová, P. Rossmann, F. Škanta, M. Bilej
The mechanism of the impact of dioxins on intestinal and chloragogenous tissue of the earthworm, Eisenia andrei, was proposed based on observations of various levels of gene expression.
- 29 Transformation and removal pathways of four common PPCP/EDCs in soil**
L.K. Dodgen, J. Li, X. Wu, Z. Lu, J.J. Gan
The majority of four PPCP/EDCs spiked into soil was mineralized or converted to bound residues. Half-lives of parents were short, but many transformation products were detected.
- 37 A systematic evaluation of agglomeration of Ag and TiO₂ nanoparticles under freshwater relevant conditions**
E. Topuz, L. Sigg, I. Talinli
Agglomeration studies of engineered nanoparticles in media simulating freshwater are useful for the prediction of their behavior in freshwater.
- 45 Stemflow acid neutralization capacity in a broadleaved deciduous forest: The role of edge effects**
A.N. Shiklomanov, D.F. Levia
Liriodendron tulipifera trees located at the forest edge have a significantly higher stemflow acid neutralization capacity than those in the forest interior at our rural study site.
- 54 Long-term source apportionment of ambient fine particulate matter (PM_{2.5}) in the Los Angeles Basin: A focus on emissions reduction from vehicular sources**
S. Hasheminassab, N. Daher, B.D. Ostro, C. Sioutas
Strict regulations have been effective in reducing PM emissions from vehicular sources in the LA Basin over the past decade.
- 65 Microplastics in bivalves cultured for human consumption**
L. Van Cauwenberghe, C.R. Janssen
*Microplastics were detected in two commercially grown bivalve species (*Mytilus edulis* and *Crassostrea gigas*), enabling us to estimate a preliminary human dietary exposure.*
- 71 Improvements in pollutant monitoring: Optimizing silicone for co-deployment with polyethylene passive sampling devices**
S.G. O'Connell, M.A. McCartney, L.B. Paulik, S.E. Allan, L.G. Tidwell, G. Wilson, K.A. Anderson
Silicone passive sampling is optimized through laboratory methods, and OPAHs highlight differences between polyethylene and silicone passive samplers in aqueous field deployments.

Continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information-Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; Thomson Scientific GeoSciTech; Science Citation Index; SciSearch). Also covered in the abstract and citation database Scopus®. Full text available on ScienceDirect®.



0269-7491(201410)193;1-G



ELSEVIER

CONTENTS—Continued from outside back cover

- 79 Effects of biochar and activated carbon amendment on maize growth and the uptake and measured availability of polycyclic aromatic hydrocarbons (PAHs) and potentially toxic elements (PTEs)**
A. Brennan, E. Moreno Jiménez, J.A. Albuquerque, C.W. Knapp, C. Switzer
Biochar and activated carbon reduce PAH and PTE uptake to maize plants to varying extents and improve plant growth.
- 88 Decreasing effect and mechanism of FeSO₄ seed particles on secondary organic aerosol in α -pinene photooxidation**
B. Chu, Y. Liu, J. Li, H. Takekawa, J. Liggio, S.-M. Li, J. Jiang, J. Hao, H. He
FeSO₄ seed particles decrease SOA mass in photooxidation of α -pinene. Free radicals are proposed to be formed in the aqueous phase and react further with the organic products.
- 94 Occurrence of antibiotics and antibiotic resistances in soils from wastewater irrigation areas in Beijing and Tianjin, China**
C. Chen, J. Li, P. Chen, R. Ding, P. Zhang, X. Li
The concentrations of antibiotics and abundances of resistance genes were significantly greater in irrigated than in non-irrigated soils in Beijing and Tianjin, China.
- 102 Remobilisation of silver and silver sulphide nanoparticles in soils**
D.A. Navarro, J.K. Kirby, M.J. McLaughlin, L. Waddington, R.S. Kookana
The potential environmental exposure to nanoparticulate Ag (Ag⁰/Ag₂S-NPs) as a result of its remobilisation in soil will be low.
- 111 Mechanistic considerations of photosensitized transformation of microcystin-LR (cyanobacterial toxin) in aqueous environments**
S. Yan, D. Zhang, W. Song
The photosensitized isomerization of MC-LR with triplet states of DOM plays a key role under solar simulated irradiation.
- 119 Tree and forest effects on air quality and human health in the United States**
D.J. Nowak, S. Hirabayashi, Allison Bodine, E. Greenfield
Air pollution modeling reveals broad-scale impacts of pollution removal by trees in urban and rural areas across the United States on air pollution concentrations and human health.
- 130 Correlation of noise levels and particulate matter concentrations near two major freeways in Los Angeles, California**
S. Shu, P. Yang, Y. Zhu
Residents live on the downwind side of freeways are exposed to higher UFP concentrations and similar noise level compared to those who live on the upwind side.
- 138 Quantifying the effects of soil temperature, moisture and sterilization on elemental mercury formation in boreal soils**
R. Pannu, S.D. Siciliano, N.J. O'Driscoll
Hg reduction kinetics in soils was studied. Cumulative Hg(0) formed and reduction rates increased with soil temperature and moisture. Biotic processes played a dominant role.
- 147 Mercury bioaccumulation in estuarine wetland fishes: Evaluating habitats and risk to coastal wildlife**
C.A. Eagles-Smith, J.T. Ackerman
Estuarine fish mercury concentrations varied substantially among wetland habitats, and managed wetlands were most important habitats for mercury risk to coastal wildlife.
- 156 Volatile organic compound identification and characterization by PCA and mapping at a high-technology science park**
C.-H. Lan, Y.-L. Huang, S.-H. Ho, C.-Y. Peng
A three-step approach, which included concentration profile estimation, principal component analysis, and concentration mapping, was used to characterize VOCs.
- 165 Impact of solar UV radiation on toxicity of ZnO nanoparticles through photocatalytic reactive oxygen species (ROS) generation and photo-induced dissolution**
H. Ma, L.K. Wallis, S. Diamond, S. Li, J. Canas-Carrell, A. Parra
Toxicity of ZnO nanoparticles is enhanced under solar UV radiation through both photocatalytic ROS generation and photo-induced dissolution.
- 173 Riverbank filtration potential of pharmaceuticals in a wastewater-impacted stream**
P.M. Bradley, L.B. Barber, J.W. Duris, W.T. Foreman, E.T. Furlong, L.E. Hubbard, K.J. Hutchinson, S.H. Keefe, D.W. Kolpin
Effluent-induced hydraulic gradients drive substantial surface-water to groundwater transport of wastewater pharmaceuticals.
- 181 Fate and metabolism of tetrabromobisphenol A in soil slurries without and with the amendment with the alkylphenol degrading bacterium *Sphingomonas* sp. strain TTNP3**
F. Li, J. Wang, P. Nastold, B. Jiang, F. Sun, A. Zenker, B.A. Kolvenbach, R. Ji, P. François-Xavier Corvini
*Degradation of TBBPA in soil slurry followed four primary pathways and was stimulated by the BPA-degrading bacterium *Sphingomonas* sp. strain TTNP3.*
- 189 Widespread occurrence of neonicotinoid insecticides in streams in a high corn and soybean producing region, USA**
M.L. Hladik, D.W. Kolpin, K.M. Kuivila
Three neonicotinoids (clothianidin, thiamethoxam, imidacloprid) were frequently detected in streams in a U.S. corn and soybean region throughout the growing season.

CONTENTS—Continued from inside back cover

- 197 Oxidative enzymatic response of white-rot fungi to single-walled carbon nanotubes**
T.D. Berry, T.R. Filley, R.A. Blanchette
Single-walled carbon nanomaterials were found to stimulate oxidative enzyme activity in cultured white-rot fungi, dependent on the purity and surface chemistry of the nanomaterials.
- 205 Root biomass production in populations of six rooted macrophytes in response to Cu exposure: Intra-specific variability versus constitutive-like tolerance**
L. Marchand, F. Nsanganwimana, J.B. Lamy, C. Quintela-Sabaris, C. Gonnelli, I. Colzi, T. Fletcher, N. Oustrière, A. Kolbas, P. Kidd, F. Bordas, P. Newell, P. Alvarenga, A. Deletic, M. Mench
Copper tolerance of four rooted macrophytes present an intra-specific variability.
- 216 Incidence of anticancer drugs in an aquatic urban system: From hospital effluents through urban wastewater to natural environment**
L. Ferrando-Ciiment, S. Rodriguez-Mozaz, D. Barceló
Integrated monitoring of 10 anticancer drugs in hospital and urban effluents as well as in receiving river waters. Assessment of their environmental impact.
- 224 Distribution of perfluoroalkyl substances (PFASs) with isomer analysis among the tissues of aquatic organisms in Taihu Lake, China**
S. Fang, S. Zhao, Y. Zhang, W. Zhong, L. Zhu
Distribution of PFASs and the isomers of PFOS and PFOA was investigated among the tissues of aquatic organisms in Taihu Lake, China.
- 233 Development of a passive sampler for Zinc(II) in urban pond waters using a polymer inclusion membrane**
M.I.G.S. Almeida, C. Chan, V.J. Pettigrove, R.W. Cattrall, S.D. Kolev
A novel passive sampler utilizing for the first time a polymer inclusion membrane as its semi-permeable barrier is proposed for monitoring Zn(II) in urban waters.
- 240 How much do PCB toxic equivalents account for PHAH toxicity in predatory birds?**
M.G. Pereira, A.J. Murk, H. Van den Berg, L.A. Walker, R.F. Shore
Significant AhR-mediated toxicity in predatory birds, at least in Britain, is most likely to be due to dioxin-like PCBs rather than other PHAHs
- 247 Explaining PAH desorption from sediments using Rock Eval analysis**
A. Poot, M.T.O. Jonker, F. Gillissen, A.A. Koelmans
Rock Eval Residual Carbon is an appropriate proxy for those condensed carbonaceous fractions in sediments that explain the magnitude of kinetic fractions for the most bioavailable PAHs.
- 254 Organophosphate flame retardants and organosiloxanes in predatory freshwater fish from locations across Canada**
D.J. McGoldrick, R.J. Letcher, E. Barresi, M.J. Keir, J. Small, M.G. Clark, E. Sverko, S.M. Backus
Of the compounds screened, 6 organophosphate flame retardants and 5 siloxanes were detected in whole body homogenate of Lake Trout or Walleye collected from Canadian water bodies.
- 262 Estimating terrestrial amphibian pesticide body burden through dermal exposure**
R.J. Van Meter, D.A. Glinski, T. Hong, M. Cyterski, W.M. Henderson, S.T. Purucker
We report pesticide body burdens in post-metamorphic amphibians after dermal exposure and are the first to identify important parameters in pesticide accumulation through predictive modeling.
- 269 Levels and profile of several classes of organic contaminants in matched indoor dust and serum samples from occupational settings of Pakistan**
N. Ali, T. Mehdi, R.N. Malik, S.A.M.A.S. Eqani, A. Kamal, A.C. Dirtu, H. Neels, A. Covaci
This study provides an important benchmark for bio-monitoring and exposure assessment of several classes of organic contaminants in human at workplaces.
- 277 Uptake and elimination kinetics of metals in soil invertebrates: A review**
M.M. Ardestani, N.M. van Straalen, C.A.M. van Gestel
Metal toxicokinetics described for soil organisms.
- 296 Evidence of widespread ozone-induced visible injury on plants in Beijing, China**
Z. Feng, J. Sun, W. Wan, E. Hu, V. Calatayud
Ozone-induced visible injury is widespread in Beijing.