





**MiniReviews**

- 1 Asymptomatic bacteriuria: prevalence rates of causal microorganisms, etiology of infection in different patient populations, and recent advances in molecular detection  
*D.S. Ipe, L. Sundac, W.H. Benjamin Jr, K.H. Moore and G.C. Ulett*
- 11 Protein-serine/threonine/tyrosine kinases in bacterial signaling and regulation  
*C. Cousin, A. Derouiche, L. Shi, Y. Pagot, S. Poncet and I. Mijakovic*

**Research Letters**

- 20 Evidence for xylooligosaccharide utilization in *Weissella* strains isolated from Indian fermented foods and vegetables  
*A. Patel, P. Falck, N. Shah, P. Immerzeel, P. Adlercreutz, H. Ståbrand, J.B. Prajapati, O. Holst and E. Nordberg Karlsson*
- 29 Evaluation of nutritional and physical stress conditions during vegetative growth on conidial production and germination in *Ophiocordyceps sinensis*  
*S.-Y. Ren and Y.-J. Yao*
- 36 Identification of two new polar tube proteins related to polar tube protein 2 in the microsporidian *Antonospora locustae*  
*V. Polonais, A. Belkorchia, M. Roussel, E. Peyretailade, P. Peyret, M. Diogon and F. Delbac*
- 45 A gene determining a new member of the SARP family contributes to transcription of genes for the synthesis of the angucycline polyketide auricin in *Streptomyces aureofaciens* CCM 3239  
*A. Rehakova, R. Novakova, L. Feckova, E. Mingyar and J. Kormanec*
- 56 Analysis of polyhydroxyalkanoate (PHA) synthase gene and PHA-producing bacteria in activated sludge that produces PHA containing 3-hydroxydodecanoate  
*C. Yang, W. Zhang, R. Liu, C. Zhang, T. Gong, Q. Li, S. Wang and C. Song*
- 65 Cortex synthesis during *Bacillus subtilis* sporulation depends on the transpeptidase activity of SpoVD  
*E. Bukowska-Faniband and L. Hederstedt*
- 73 Transport of the botulinum neurotoxin-associating protein, nontoxic nonhemagglutinin, across the rat small intestinal epithelial cell monolayer  
*S.-I. Miyashita, Y. Sagane, K. Niwa and T. Watanabe*