


T14
F40

Volume 337, January 15 2013

ISSN 0378-3812



FLUID PHASE

EQUILIBRIA

AN INTERNATIONAL JOURNAL

FLUID PHASE

EQUILIBRIA

CONTENTS

(Abstracted/Indexed in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, SciVerse Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®*)

Full Length Articles

- Partitioning of alkaline protease from *Bacillus licheniformis* (ATCC 21424) using PEG-K₂HPO₄ aqueous two-phase system
M. Yavari, G.R. Pazuki, M. Vossoughi, S.A. Mirkhani and A.A. Seifkordi (Tehran, Iran) 1–5
- Comparison between three predictive methods for the calculation of polymer solubility parameters
I. Díaz, E. Díez, J. Camacho, S. León and G. Ovejero (Madrid, Spain) 6–10
- Ionic liquids as solvents to separate the azeotropic mixture hexane/ethanol
S. Corderí, B. González (Vigo, Spain), N. Calvar (Porto, Portugal) and E. Gómez (Vigo, Spain) 11–17
- Physical-chemical parameters promoting phase changes in non-ionic environmental-friendly microemulsions
M.C. Fleancu, N.L. Olteanu, A.E. Rogozea, A.V. Crisciu, I. Pincovschi and M. Mihaly (Bucharest, Romania) 18–25
- A study on thermodynamics effect of [EMIM]-Cl and [OH-C₂MIM]-Cl on methane hydrate equilibrium line
B. Partoon, N.M.S. Wong, K.M. Sabil, K. Nasrifar and M.R. Ahmad (Tronoh, Malaysia) 26–31
- Experimental study of phase equilibria in aqueous mixtures of phosphoric acid with isoamyl acetate and methyl isoamyl ketone at $T = (298.2, 308.2, \text{ and } 318.2) \text{ K}$
H. Ghanadzadeh Gilani, A. Ghanadzadeh Gilani and S. Shekarsaraee (Rasht, Iran) 32–38
- Physico-chemical studies of oppositely charged protein-surfactant system in aqueous solutions: Sodium dodecyl sulphate (SDS)-lysozyme
S. Chauhan, M.S. Chauhan, P. Sharma, D.S. Rana (Shimla, India) and A. Umar (Najran, Saudi Arabia) 39–46
- Liquid-liquid extraction of toluene from heptane by {[4bmpy][Tf₂N] + [emim][CHF₂CF₂SO₃]} ionic liquid mixed solvents
S. García, J. García, M. Larriba, A. Casas and F. Rodríguez (Madrid, Spain) 47–52
- Prediction of vapor pressure and heats of vaporization of edible oil/fat compounds by group contribution
R. Ceriani (Campinas, Brazil), R. Gani (Lyngby, Denmark) and Y.A. Liu (Blacksburg, VA, USA) 53–59
- Solubility of hydrofluorocarbons in aromatic solvents and alcohols: Experimental data and modeling with CPA EoS
J.M.M.V. Sousa (Coimbra, Portugal), A.J. Queimada, E.A. Macedo (Porto, Portugal) and I.M.A. Fonseca (Coimbra, Portugal) 60–66
- Liquid-liquid equilibria of imidazolium ionic liquids having bistriflamide or triflate anions with aromatic non-polar and polar compounds
M.S. Calado (Lisboa, Portugal), Z. Petrovski, M.S. Manic, V. Najdanovic-Visak (Caparica, Portugal), E.A. Macedo (Porto, Portugal) and Z.P. Visak (Lisboa, Portugal) 67–72
- A simple correlation for prediction of heat capacities of ionic liquids
N. Farahani, F. Gharagheizi, S.A. Mirkhani (Buinzhahra, Iran) and K. Tumba (Durban, South Africa) 73–82
- Experiment and model for the surface tension of carbonated MEA-MDEA aqueous solutions
D. Fu, L. Wei and S. Liu (Baoding, China) 83–88
- Asphaltene deposition prediction using adaptive neuro-fuzzy models based on laboratory measurements
K. Salahshoor, S. Zakeri, S. Mahdavi, R. Kharrat and M. Khalifeh (Tehran, Iran) 89–99
- Correlating and predicting the solubilities of polycyclic aromatic hydrocarbons in supercritical fluids using the compressed gas model and the reference solubilities
B. Shen, L. Feng, S. Li and H. Li (Tianjin, PR China) 100–108
- Isobaric vapor-liquid equilibrium of binary systems of decane with p-, o-, m-xylene at 20 kPa
S. Yang, Y. Wang, X. Wang, Y. Zhu and N. Chen (Xinxiang, PR China) 109–114
- Phase equilibrium of ionic semicathrate hydrates formed with tetrabutylammonium bromide and tetrabutylammonium chloride
K. Sato, H. Tokutomi and R. Ohmura (Yokohama, Japan) 115–118
- Thermal conductivity, rheological behaviour and density of non-Newtonian ethylene glycol-based SnO₂ nanofluids
A. Mariano (Neuquén, Argentina), M.J. Pastoriza-Gallego, L. Lugo (Vigo, Spain), A. Camacho, S. Canzonieri (Neuquén, Argentina) and M.M. Piñeiro (Vigo, Spain) 119–124
- Thermodynamics of alkanone + aromatic hydrocarbon mixtures
J.A. González, I. Alonso (Valladolid, Spain), C. Alonso-Tristán (Burgos, Spain), I.G. De La Fuente and J.C. Cobos (Valladolid, Spain) 125–136
- Swelling equilibrium of hydrogels of (N-isopropyl acrylamide + anionic and cationic comonomers) in aqueous solutions of sodium chloride: Experimental results and modeling
L. Ninni, V. Ermatchkov, H. Hasse and G. Maurer (Kaiserslautern, Germany) 137–149

(Contents continued on BM I)



(contents continued)

Clathrate hydrate equilibria in light olefins and mixed methane-olefins systems A. Chapoy, I. Alsiyabi, J. Gholinezhad, R. Burgass and B. Tohidi (Edinburgh, UK).....	150–155
Volumetric behavior of the ternary system benzene-2-methoxy-2-methylbutane-2,2,4-trimethylpentane and all binary sub-systems at temperature range (298.15–318.15) K H. Houšková, L. Morávková, Z. Sedláková (Prague, Czech Republic), T. Boublík and Z. Kolská (Labem, Czech Republic).....	156–164
Concentration dependent surface area parameter model for calculation of activity coefficients Y. Iwai and Y. Yamamoto (Fukuoka, Japan).....	165–173
Simple isotherm equations to fit type I adsorption data M.A. Mosquera (West Lafayette, IN, USA).....	174–182
Simple yet accurate prediction of liquid molar volume via their molecular structure M. Bagheri (Tehran, Iran), T.N.G. Borhani and G. Zahedi (Johor, Malaysia).....	183–190
Cuckoo Search: A new nature-inspired optimization method for phase equilibrium calculations V. Bhargava (Kharagpur, India), S.E.K. Fateen (Egypt) and A. Bonilla-Petriciolet (Aguascalientes, Mexico).....	191–200
A practical method for the calculation of liquid-liquid equilibria in multicomponent organic-water-electrolyte systems using physicochemical constraints A. Zuend and J.H. Seinfeld (Pasadena, CA, USA).....	201–213
New volume translated PR equation of state for pure compounds and gas condensate systems M. Nazarzadeh and M. Moshfeghian (Shiraz, Iran and Norman, OK, USA).....	214–223
Phase diagrams for liquid-liquid and liquid-solid equilibrium of the ternary polyethylene glycol + di-sodium hydrogen citrate + water system M.T. Zafarani-Moattar and P. Jafari (Tabriz, Iran).....	224–233
Vapour-liquid equilibrium (VLE) for the systems furan + n-hexane and furan + toluene. Measurements, data treatment and modeling using molecular models M. Nala (Durban, South Africa and Fontainebleau, France), E. Auger (Villetaneuse, France), I. Gedik, N. Ferrando (Rueil-Malmaison, France), M. Dicko (Villetaneuse, France), P. Paricaud (Palaiseau, France), F. Volle, J.P. Passarello (Villetaneuse, France), J.-C. de Hemptinne (Rueil-Malmaison, France), P. Tobaly (Villetaneuse, France), P. Stringari, C. Coquelet (Fontainebleau, France), D. Ramjugernath, P. Naidoo (Durban, South Africa) and R. Lugo (Rueil-Malmaison, France).....	234–245
Measurement and correlation of saturated liquid properties and gas solubility for decane, tetradecane and their binary mixtures saturated with carbon dioxide H. Nourozieh, M. Kariznovi and J. Abedi (Calgary, Canada).....	246–254
Excess molar enthalpies for binary mixtures of 1,2-dichloroethane with ethers at 298.15 K and atmospheric pressure F. Amireche-Ziar (Alger, Algeria), D. Richon (Fontainebleau, France) and F.B. Belaribi (Alger, Algeria).....	255–258
On the fundamental derivative of gas dynamics in the vapor-liquid critical region of single-component typical fluids N.R. Nannan (Paramaribo, Suriname and Delft, The Netherlands), A. Guardone (Milano, Italy) and P. Colonna (Delft, The Netherlands).....	259–273
Application of the statistical associating fluid theory for potentials of variable range (SAFT-VR) coupled with renormalisation-group (RG) theory to model the phase equilibria and second-derivative properties of pure fluids E. Forte (London, UK), F. Llovel (Barcelona, Spain), J.P.M. Trusler and A. Galindo (London, UK).....	274–287
Prediction of the solubility of quartz in salt solutions from 25 °C to 900 °C using the 3-parameter Non-Random Two-Liquid (NRTL) model A.G. Carr and J.W. Tester (Ithaca, NY, USA).....	288–297
Distribution of MEG and methanol in well-defined hydrocarbon and water systems: Experimental measurement and modeling using the CPA EoS M. Riaz, M.A. Yussuf, G.M. Kontogeorgis, E.H. Stenby, W. Yan (Lyngby, Denmark) and E. Solbraa (Trondheim, Norway).....	298–310
A revised LIQUAC and LIFAC model (LIQUAC*/LIFAC*) for the prediction of properties of electrolyte containing solutions A. Mohs and J. Gmehling (Oldenburg, Germany).....	311–322
Experimental study of the PVT and C_VVT properties of <i>n</i> -butanol in the critical region S.M. Rasulov, L.M. Radzhabova, I.M. Abdulagatov and G.V. Stepanov (Dagestan, Russia).....	323–353
Thermodynamic analysis and correlation of solubility of candesartan cilexetil in aqueous solvent mixtures P. Cui, Q. Yin, J. Gong, Y. Wang, H. Hao, C. Xie, Y. Bao, M. Zhang, B. Hou and J. Wang (Tianjin, People's Republic of China).....	354–362
Phase behavior and compressibility factor of two China gas condensate samples at pressures up to 95 MPa H. Liu, C.-Y. Sun, K.-L. Yan, Q.-L. Ma, J. Wang, G.-J. Chen (Beijing, China), X.-J. Xiao, H.-Y. Wang (Korla, China), X.-T. Zheng and S. Li (Beijing, China).....	363–369
Adsorption and micellar phase properties of anionic surfactant in the presence of electrolyte and oil at different temperatures P. Moradi, B. Sohrabi, M. Najafi and V. Khani (Tehran, Iran).....	370–378
Short Communication	
On the description of the <i>liquidus</i> line of systems presenting peritectic reactions D.F. Barbosa and P. de Alcântara Pessôa Filho (São Paulo, Brazil).....	379–383