


Volume 338, January 25 2013

ISSN 0378-3812



FLUID PHASE

EQUILIBRIA

AN INTERNATIONAL JOURNAL

FLUID PHASE

EQUILIBRIA

CONTENTS

(Abstracted/Indexed in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, SciVerse Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®*)

Full Length Articles

- Evaluation of the PC-SAFT, SAFT and CPA equations of state in predicting derivative properties of selected non-polar and hydrogen-bonding compounds
A.J. de Villiers, C.E. Schwarz, A.J. Burger (Matieland, South Africa) and G.M. Kontogeorgis (Lyngby, Denmark) 1–15
- Isothermal vapor-liquid equilibria and excess Gibbs free energies in some binary nitroalkane + chloroalkane mixtures at temperatures from 298.15 K to 318.15 K
D. Dragoescu, M. Teodorescu and D. Gheorghe (Bucharest, Romania) 16–22
- Reliability of the correlation allowing the k_{ij} to switch from an alpha function to another one in hydrogen-containing systems
J.-N. Jaubert, J. Qian, R. Privat (Nancy, France) and C.F. Leibovici (Pau, France) 23–29
- Vapor-liquid equilibrium for the ternary carbon dioxide-ethanol-nonane and decane systems
M.G. Arenas-Quevedo, L.A. Galicia-Luna, O. Elizalde-Solis (México, Mexico) and J.A. Pérez-Pimienta (Tepic, Mexico) 30–36
- The study of phase behavior of aqueous 1-propanol/2-propanol/2-butanol/2-methyl-2-propanol systems in the presence of disodium tartrate or disodium succinate at $T = 298.15$ K
M.T. Zafarani-Moattar, V. Hosseinpour-Hashemi (Tabriz, Iran), S. Banisaeid and M.A.S. Beirami (Urmia, Iran) 37–45
- Use of a hybrid optimization method to reduce vapor-liquid equilibrium data of maverick systems: The case of carbon dioxide with 2-methoxyethanol and 2-ethoxyethanol using cubic equations of state
B. Ebrahimi, T. Hatami (Sanandaj, Iran) and J.H. Vera (Montreal, Canada) 46–53
- A Pitzer-based characterization of aqueous magnesium chloride, calcium chloride and potassium iodide solution densities to high temperature and pressure
D. Rowland and P.M. May (Murdoch, Australia) 54–62
- MOQUAC, a new expression for the excess Gibbs energy based on molecular orientations
R. Bronneberg (Aachen, Germany) and A. Pfennig (Graz, Austria) 63–77
- Experimental data for liquid-liquid equilibrium of fatty systems with emphasis on the distribution of tocopherols and tocotrienols
M. Ansolin, R.C. Basso, A.J. de Almeida Meirelles and E.A.C. Batista (Campinas, Brazil) 78–86
- Vapor-liquid-liquid-hydrate phase equilibrium calculation for multicomponent systems containing hydrogen
Q.-L. Ma, G.-J. Chen and C.-Y. Sun (Beijing, PR China) 87–94
- Vapour-liquid equilibria and excess enthalpies of the binary mixtures 1-pentanol with 2,2,4-trimethylpentane or *n*-heptane
A. Moreau, M.C. Martín (Valladolid, Spain), F. Aguilar (Burgos, Spain) and J.J. Segovia (Valladolid, Spain) 95–99
- Phase equilibria with hydrate formation in $H_2O + CO_2$ mixtures modeled with reference equations of state
A. Jäger (Bochum, Germany), V. Vinš (Prague, Czech Republic), J. Gernert, R. Span (Bochum, Germany) and J. Hrubý (Prague, Czech Republic) 100–113
- Measurements for the dissociation conditions of methane hydrate in the presence of 2,5-dihydrofuran and 3,4-dihydro-2H-pyran
C.-T. Cheng, L.-J. Chen, S.-T. Lin, M. Tang (Taipei, Taiwan), P.-C. Chen (New Taipei City, Taiwan) and Y.-P. Chen (Taipei, Taiwan) 114–118
- A correlation to predict speed of sound in liquids: 1. *n*-Alkanes ($\geq C_3$) and their mixtures at high pressures
H. Padilla-Victoria, G.A. Iglesias-Silva (Celaya, Mexico), M. Ramos-Estrada (Morelia, Mexico) and K.R. Hall (College Station, TX, USA) 119–127
- Vapor-liquid equilibrium of the ethanol + 3-methyl-1-butanol system at 50.66, 101.33 and 151.99 kPa
J.A. Duran, F.P. Córdoba, I.D. Gil, G. Rodríguez and A. Orjuela (Bogotá, Colombia) 128–134
- Prediction of different thermodynamic properties for systems of alcohols and sulfate-based anion Ionic Liquids using modified UNIFAC
T. Hector, L. Uhlig and J. Gmehling (Oldenburg, Germany) 135–140
- Experimental and VTPR-predicted volumetric properties of branched hexanes
H. Guerrero, M. García-Mardones, V. Pérez-Gregorio, I. Gascón and C. Lafuente (Zaragoza, Spain) 141–147
- Vapor pressures, standard molar enthalpies, entropies Gibbs energies of sublimation and heat capacities of 2,5- and 3,5-dibromobenzoic acids
S. Vecchio and B. Brunetti (Rome, Italy) 148–154
- Near-infrared spectroscopic investigation of water in supercritical CO_2 and the effect of $CaCl_2$
Z. Wang, A.R. Felmy, C.J. Thompson, J.S. Loring, A.G. Joly, K.M. Rosso, H.T. Schaefer (Richland, WA, USA) and D.A. Dixon (Tuscaloosa, AL, USA) 155–163

(Contents continued on BM I)



(contents continued)

Measurements of H ₂ S solubility in aqueous diisopropanolamine solutions and vapour pressure of diisopropanolamine P. Uusi-Kyyny, C. Dell'Era, A. Penttilä, M. Pakkanen and V. Alopaeus (Aalto, Finland)	164–171
Solid-liquid phase diagram of the two Ionic Liquids EMIMCl and BMIMCl M. Kick, P. Keil and A. König (Erlangen, Germany)	172–178
Modeling of hydrate formation conditions for CH ₄ , C ₂ H ₆ , C ₃ H ₈ , N ₂ , CO ₂ and their mixtures using the PRSV2 equation of state and obtaining the Kihara potential parameters for these components G. Moradi (Kermanshah, Iran) and E. Khosravani (Tehran, Iran)	179–187
A novel static analytical apparatus for phase equilibrium measurements C. Narasigadu, P. Naidoo (Durban, South Africa), C. Coquelet (Durban, South Africa and Fontainebleau, France), D. Richon and D. Ramjugernath (Durban, South Africa)	188–196
Salt effects on the diffusion of caffeine in aqueous solutions A.C.F. Ribeiro, C.I.A.V. Santos, M.C.F. Barros, J.C.S. Gomes, L.M.P. Veríssimo, M. da G. Miguel (Coimbra, Portugal), M.A. Esteso (Madrid, Spain), P. Lo Nostro (Florence, Italy), V.M.M. Lobo (Coimbra, Portugal) and D.G. Leaist (Nova Scotia, Canada)	197–203
Isoopiestic studies of thermodynamic properties of solutions of ampicillin sodium and penicillin sodium in water at $T = 298.15$ K J. Azamat, S.M.S. Ahmadian, J.J. Sardroodi and M. Fouladian (Tabriz, Iran)	204–208
Ammonium ionic liquids as green solvents for drugs C.I. Melo (Caparica, Portugal), R. Bogel-Lukasik (Lisboa, Portugal), M. Nunes da Ponte and E. Bogel-Lukasik (Caparica, Portugal)	209–216
Determination and modeling of aqueous solubility of 4-position substituted benzoic acid compounds in a high-temperature solution N. Sunsandee, S. Suren (Bangkok, Thailand and Bratislava, Slovak Republic), N. Leepipatpiboon (Bangkok, Thailand), M. Hronec (Bratislava, Slovak Republic) and U. Pancharoen (Bangkok, Thailand)	217–223
The choice of solvent and liquid-liquid equilibrium for ternary water + 2-methylaziridine + chloroform system: Experimental data and modeling J. Saien, M. Norouzi and H. Dehghani (Hamedan, Iran)	224–231
Steam solubilities of solid MoO ₃ , ZnO and Cu ₂ O, calculated on a basis of a thermodynamic model N.N. Akinfiev (Moscow, Russia) and A.V. Plyasunov (Moscow Region, Russia)	232–244
Fourth order virial equation of state for spherical molecules using semi-soft core potential function S.J. Pai and Y.C. Bae (Seoul, Republic of Korea)	245–252
Ternary liquid-liquid equilibria of trifluorotris(perfluoroethyl)phosphate based ionic liquids + methanol + heptane M. Wlazło and A. Marciniak (Warsaw, Poland)	253–256
The role of thermal path on the accuracy of gas hydrate phase equilibrium data using isochoric method M. Mohammad-Taheri, A. Zarringhalam Moghaddam, K. Nazari (Tehran, Iran) and N. Gholipour Zanjani (Tehran, Iran and Karaj, Iran)	257–264
Solid-liquid equilibrium of 1-hydroxyethane-1,1-diphosphonic acid in different solvents X. Zhang, X. Yang, L. Hao and H. Wei (Tianjin, China)	265–268
Generalized expression of chemical potential with influence of external fields and its applications: Effect of charged particles on droplet condensation G. Han and H. Wang (Guangdong, China)	269–273
Thermodynamic study of electrolyte mixture (KCl + KNO ₃ + water + proline) by potentiometric measurements at $T = 298.2$ and 303.2 K B. Ghalami-Chooabar and F. Sayyadi-Nodehi (Rasht, Iran)	274–281

Short Communication

Experimental measurement and modelling of viscosity of the binary systems pyridine or nicotine with polyethylene glycols at $T = (288.15-333.15)$ K. New UNIFAC-VISCO and ASOG-VISCO interaction parameters D.M. Bajić, J. Jovanović, E.M. Živković (Belgrade, Serbia), Z.P. Visak (Lisboa, Portugal), S.P. Šerbanović and M. Lj. Kijevčanin (Belgrade, Serbia)	282–293
--	---------