

ПИ

F70



ELSEVIER

Volume 341, March 15 2013

ISSN 0378-3812

FLUID PHASE EQUILIBRIA

AN INTERNATIONAL JOURNAL

FLUID PHASE EQUILIBRIA

CONTENTS

(Abstracted/Index in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, SciVerse Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®*)

Full Length Articles

Modeling the melting temperature depression of ionic liquids caused by supercritical carbon dioxide J.O. Valderrama (La Serena, Chile) and P.F. Arce (Campinas, Brazil)	1–6
Solubility of succinic acid in different aqueous solvent mixtures: Experimental measurement and thermodynamic modeling X. Jiang, Y. Hu, Z. Meng, W. Yang and F. Shen (Nanjing, China)	7–11
Thermodynamic simulation of transesterification reaction by Gibbs energy minimization D.M. Yancy-Caballero and R. Guirardello (Campinas, Brazil)	12–22
Phase behavior of aqueous two-phase systems composed of 1-ethyl-3-methylimidazolium tetrafluoroborate and phosphate-based salts at different temperatures H. Lv, D. Guo, Z. Jiang, Y. Li and B. Ren (Zhengzhou, PR China)	23–29
Investigation on vapor-liquid equilibrium for 2-propanol + 1-butanol + 1-pentanol at 101.3 kPa J. Wang and Z. Bao (Nanjing, PR China)	30–34
Ternary (liquid-liquid) equilibria of nitrate based ionic liquid + alkane + benzene at 298.15 K: Experiments and correlation B. Mokhtarani (Tehran, Iran), J. Musavi, M. Parvini (Semnan, Iran), M. Mafi, A. Sharifi and M. Mirzaei (Tehran, Iran)	35–41
Study on the effect of solvent on the inclusion interaction of hydroxypropyl- β -cyclodextrin with three kinds of coumarins by phase solubility method A. Chen, M. Liu, L. Dong and D. Sun (Liaocheng, People's Republic of China)	42–47
High-temperature vapour-liquid equilibrium for the water-alcohol systems and modeling with SAFT-VR: 1. Water-ethanol A.F. Cristina, S. Rosa, P. Morgado (Lisboa, Portugal), A. Galindo (London, UK), E.J.M. Filipe, A.M.F. Palavra and C.A. Nieto de Castro (Lisboa, Portugal)	48–53
Phase equilibria for the ternary systems ethanol, water + ethylene glycol or + glycerol at 101.3 kPa J. Pla-Franco, E. Lladosa, S. Loras and J.B. Montón (Valencia, Spain)	54–60
Experiment and thermodynamic modeling of methane hydrate equilibria in the presence of aqueous imidazolium-based ionic liquid solutions using electrolyte cubic square well equation of state M. Zare, A. Haghtalab, A.N. Ahmadi and K. Nazari (Tehran, Iran)	61–69
Liquid-liquid equilibria of aqueous two-phase systems of the ionic liquid brominated N-ethyl pyridine and sodium dihydrogen phosphate, sodium sulfate, ammonium citrate, and potassium tartrate at different temperatures: Experimental determination and correlation Y.-L. Li, M.-S. Zhang, H. Su, Q. Liu and W.-S. Guan (Xi'an, China)	70–77
An equation of state for the thermodynamic properties of 1,1,1,2,3-hexafluoropropane (R236ea) X. Rui, J. Pan and Y. Wang (Hangzhou, China)	78–85
Solubility of sulfapyridine in propylene glycol + water mixtures and correlation with the Jouyban-Acree model D.R. Delgado, G.A. Rodríguez, A.R. Holguín, F. Martínez (Bogotá, Colombia) and A. Jouyban (Tabriz, Iran)	86–95
Calculation of liquid-liquid equilibrium in polymer electrolyte solutions using PHSC-electrolyte equation of state M. Valavi, M.R. Dehghani and F. Feyzi (Tehran, Iran)	96–104
Multiproperty modeling for a set of binary systems. Evaluation of a model to correlate simultaneously several mixing properties of methyl ethanoate + alkanes and new experimental data L. Fernández, E. Pérez, J. Ortega (Canary Islands, Spain), J. Canosa (Spain) and J. Wisniak (Beer-Sheva, Israel)	105–123

