


NY

F 70



Volume 352, August 25 2013

ISSN 0378-3812

# **FLUID PHASE EQUILIBRIA**

**AN INTERNATIONAL JOURNAL**

# **FLUID PHASE EQUILIBRIA**

## CONTENTS

(Abstracted/Indexed in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, SciVerse Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®*)

**Articles**

Thermodynamics of solubility of thiomalic acid in different organic solvents from 278.15 K to 333.15 K Z. Meng, Y. Hu, Y. Kai, W. Yang, Z. Cao and F. Shen (Nanjing, China) .....	1–6
Study of solvation consequences of $\alpha$ -amino acids in aqueous ionic liquid solution probed by physicochemical approach M.N. Roy, P. De and P.S. Sikdar (Darjeeling, India) .....	7–13
Correlation of solubility of pioglitazone hydrochloride in different binary solvents M. Tao (Tianjin, People's Republic of China), H. Sun (Hebei, People's Republic of China), Z. Wang, P. Cui and J. Wang (Tianjin, People's Republic of China) .....	14–21
Effect of solvent on the volumetric behavior of <i>N,N'</i> -salicylidenephenyl diamine (Salophen) Schiff base at different temperatures (288.15–318.15) K R. Elhami-Kalvanagh (Mamaghan, Iran), H. Shekaari (Tabriz, Iran) and A. Kazempour (Mamaghan, Iran) .....	22–27
Conductivity in $MnSO_4$ -saccharide-water solutions at 298.15 K J. Zhao, Y. Chen, Y. Liu and K. Zhuo (Henan, People's Republic of China) .....	28–33
Correlating bubble points of ternary systems involving nine solvents and two ionic liquids using artificial neural network A.Z. Hezave, M. Lashkarbolooki and S. Raeissi (Shiraz, Iran) .....	34–41
Effect of hydrophilic ionic liquid on aggregation behavior of aqueous solutions of sodium dodecylsulfate (SDS) A. Pal and S. Chaudhary (Kurukshetra, India) .....	42–46
Isobaric vapor-liquid equilibrium for methanol + dimethyl carbonate + phosphoric-based ionic liquids F. Cai, X. Wu, C. Chen, X. Chen, C. Asumana, M.R. Haque and G. Yu (Beijing, PR China) .....	47–53
Empirical correlation of the surface tension versus the viscosity for saturated normal liquids X. Li (Qufu, PR China), J. Tian (Qufu, PR China and Dalian, PR China) and A. Mulero (Badajoz, Spain) .....	54–63
Comment on Cuckoo search: A new nature-inspired optimization method for phase equilibrium calculations by V. Bhargava, S. Fateen, A. Bonilla-Petriciolet S. Walton, M.R. Brown, O. Hassan and K. Morgan (Wales, UK) .....	64–66
$CO_2$ solubility measurement in 1-hexyl-3-methylimidazolium ([HMIM]) cation based ionic liquids J.-H. Yim and J.S. Lim (Seoul, Republic of Korea) .....	67–74
Isobaric VLE data for the system of butan-1-ol + butyl ethanoate + 1-butyl-3-methylimidazolium bis[(trifluoromethyl)sulfonyl]imide Z. Tian, X. Cui, J. Cai, Y. Zhang, T. Feng, Y. Peng and L. Xue (Tianjin, China) .....	75–79
Solubility of ethylene in norbornene + toluene + cyclic olefin copolymer systems Y. Sato, N. Hosaka, H. Inomata (Sendai, Japan) and K. Kanaka (Fuji, Japan) .....	80–85
Isobaric vapor-liquid equilibrium for binary systems of toluene + o-xylene, benzene + o-xylene, nonane + benzene and nonane + heptane at 101.3 kPa B.S. Gupta and M.-J. Lee (Taipei, Taiwan) .....	86–92
Comments on the modeling of hydrogen and hydrogen-containing mixtures with cubic equations of state U.K. Deiters (Köln, Germany) .....	93–96
Vapor-liquid equilibria of binary mixtures of propylene oxide with either ethyl benzene, 2-methylpentane, or 2-methyl-1-pentene S. Raeissi (Shiraz, Iran), L.J. Florusse (Delft, The Netherlands) and C.J. Peters (Abu Dhabi, United Arab Emirates and Eindhoven, The Netherlands) .....	97–99
Environmentally friendly solutions of liquid poly(ethylene glycol) and imidazolium based ionic liquids with bistriflamide and triflate anions: Volumetric and viscosity studies J.M. Vuksanovic (Belgrade, Serbia), M.S. Calado (Lisboa, Portugal), G.R. Ivanis, M.Lj. Kijevcanin, S.P. Serbanovic (Belgrade, Serbia) and Z.P. Visak (Lisboa, Portugal) .....	100–109
Solubility of citric acid in water, ethanol, n-propanol and in mixtures of ethanol + water M.L.N. Oliveira (Rondonopolis, Brazil), R.A. Malagoni and M.R. Franco (Uberlândia, Brazil) .....	110–113
Prediction of vapor-liquid equilibria of binary systems containing esters by using Wilson equation with parameters estimated from pure-component properties S. Kobuchi, K. Takakura (Ube-shi, Japan), S. Yonezawa (Fukuoka, Japan), K. Fukuchi (Ube-shi, Japan) and Y. Arai (Fukuoka, Japan) .....	114–117

