



Volume 360, December 25 2013

ISSN 0378-3812

# **FLUID PHASE EQUILIBRIA**

**AN INTERNATIONAL JOURNAL**

**FLUID PHASE  
EQUILIBRIA**

## CONTENTS

(Abstracted/Indexed in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA*, Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®)

Investigation on vapor–liquid equilibria for binary systems of metal ion-containing ionic liquid [bmim]Zn <sub>2</sub> Cl <sub>5</sub> /NH <sub>3</sub> by experiment and modified UNIFAC model W. Chen, S. Liang, Y. Guo, X. Gui and D. Tang (Beijing, China) .....	1–6
Estimation physicochemical property of ionic liquid [Smim][HSO <sub>4</sub> ] by semi-empirical method H. Wang, L.-j. Yang, Y.-q. Yang, D.-W. Fang and S.-l. Zang (Shenyang, PR China) .....	7–9
Temperature effect on the adsorption equilibrium at the silica–polyethylene glycol solution interface M. Wiśniewska, K. Szewczuk-Karpisz and I. Ostolska (Lublin, Poland) .....	10–15
Diffusivity data and reaction kinetics as probes for self-assembling: Cationic surfactant–chloroform system E.P. Zhiltsova, V.V. Syakaev, S.S. Lukashenko, A.P. Timosheva, R.R. Kashapov, S.K. Latypov, L.Ya. Zakharova and A.I. Konovalov (Russian Federation) .....	16–22
Sorption of CO <sub>2</sub> , C <sub>2</sub> H <sub>2</sub> and C <sub>2</sub> H <sub>4</sub> in HOF-1a studied by PHSC equation of state F. Sabzi and H. Molaei (Shiraz, Iran) .....	23–28
A group contribution method for the influence of the temperature in the viscosity of magnetic ionic liquids C.I. Daniel (Caparica, Portugal), J. Albo, E. Santos (Santander, Spain), C.A.M. Portugal, J.G. Crespo (Caparica, Portugal) and A. Irabien (Santander, Spain) .....	29–35
The effect of aqueous alcohols (methanol, t-butanol) and sulfolane on the dissociation constants and thermodynamic properties of alkanolamines N. Ramachandran, E.S. Hamborg (AH Enschede, The Netherlands) and G.F. Versteeg (AB Groningen, The Netherlands) .....	36–43
Accurate determination of thermodynamic properties for liquid alloys based on <i>ab initio</i> molecular dynamics simulation J. Han (Xiamen, China), W.Y. Wang (University Park, PA, USA and Beijing, China), C. Wang (Xiamen, China), Y. Wang (University Park, PA, USA), X. Liu (Xiamen, China) and Z.-K. Liu (University Park, PA, USA) .....	44–53
Liquid–liquid equilibrium data for the systems {LTTM + benzene + hexane} and {LTTM + ethyl acetate + hexane} at different temperatures and atmospheric pressure A.S.B. Gonzalez, M. Francisco, G. Jimeno (Eindhoven, Netherlands), S.L.G. de Dios (Santiago de Compostela, Spain) and M.C. Kroon (Eindhoven, Netherlands) .....	54–62
Study on the thermodynamic properties for ionic liquid [C <sub>4</sub> mim][OAc](1-butyl-3-methylimidazolium acetate) W. Guan, Q.-B. Zhang, X.-X. Ma, J. Wei, Y. Pan and J.-Z. Yang (Shenyang, PR China) .....	63–67
An assessment test for phase equilibrium data of water soluble and insoluble clathrate hydrate formers P. Ilani-Kashkouli, S. Babaee, F. Gharagheizi, H. Hashemi (Durban, South Africa), A.H. Mohammadi (Durban, South Africa and Paris, France) and D. Ramjugernath (Durban, South Africa) .....	68–76
Optimum extraction equilibria of the systems (water + carboxylic acid + 1-hexanol/Alamine): Thermodynamic modeling A. Senol (Istanbul, Turkey) .....	77–87
Solubility and solution thermodynamics of sulfamerazine and sulfamethazine in some ethanol + water mixtures D.R. Delgado and F. Martínez (Bogotá D.C., Colombia) .....	88–96
Solubility of 1,3,2-dioxaphosphorinane-5,5-dimethyl-2-(2,4,6-tribromophenoxy)-2-oxide in selected solvents G.-Q. Sun, L.-S. Wang, J.-C. Du and C.-M. Qi (Beijing, PR China) .....	97–105
Vapor pressures of dimethylcadmium, trimethylbismuth, and tris(dimethylamino)antimony P. Morávek, M. Fulém, J. Pangrác, E. Hulicius and K. Růžička (Prague 6, Czech Republic) .....	106–110
Vapor–liquid equilibria of binary systems composed of polyoxyethylene 4-octylphenyl ether and alcohols: Experimental measurements and correlation I. Khoiroh and M.-J. Lee (Taipei, Taiwan) .....	111–117
Measurement and correlation of the solubility of florfenicol in binary 1,2-propanediol + water mixtures from 293.15 K to 316.25 K J. Zhou, H. Fu, H. Cao, C. Lu, C. Jin, T. Zhou, M. Liu and Y. Zhang (Yaan, People's Republic of China) .....	118–123
PVTx measurements of the carbon dioxide + <i>trans</i> -1,3,3,3-tetrafluoroprop-1-ene binary system G. Di Nicola, G. Passerini, F. Polonara (Via Brecce Bianche, Italy) and R. Stryjek (Warsaw, Poland) .....	124–128
High pressure phase behavior of the binary system (ethyl lactate + carbon dioxide) A.B. Paninho (Oeiras, Portugal), A.V.M. Nunes, A. Paiva (Caparica, Portugal) and V. Najdanovic-Visak (Lancaster, UK) ....	129–133
Solubility of erythritol in methanol, ethanol and methanol + ethanol: Experimental measurement and thermodynamic modeling X. Liu, Y. Hu, W. Yang, Y. Liu and M. Liang (Nanjing, China) .....	134–138
Effect of different families (imidazolium and pyridinium) of ionic liquids-based surfactants on interfacial tension of water/crude oil system A.Z. Hezave, S. Dorostkar, S. Ayatollahi, M. Nabipour and B. Hemmateenejad (Shiraz, Iran) .....	139–145

(Contents continued on BM I)



(contents continued)

Prediction of stability limits and critical conditions of binary liquid mixtures containing nitro-methane using the NRTL model M. Qasim and N.A. Darwish (Sharjah, United Arab Emirates).....	146–155
Solubility of glycine in binary system of ethanol + water solvent mixtures: Experimental data and thermodynamic modeling Z. Cao, Y. Hu, J. Li, Y. Kai and W. Yang (Nanjing, China).....	156–160
Gas hydrate phase equilibrium in porous media: An assessment test for experimental data P. Ilani-Kashkouli, H. Hashemi, F. Gharagheizi, S. Babaee (Durban, South Africa), A.H. Mohammadi (Durban, South Africa and Paris, France) and D. Ramjugernath (Durban, South Africa).....	161–168
Structures and hydrogen bonds of biodegradable naphthenate ionic liquids H. He, S. Zhang, X. Liu, J. Wang, X. Yao and X. Zhang (Beijing, China).....	169–179
Spectroscopic studies of interaction of safranine T with ionic surfactants S. Mondal, B. Doloi and S. Ghosh (Kolkata, India).....	180–187
Concentration dependence of single-ion activity coefficients: An analysis D.P. Zarubin (Moscow, Russian Federation) .....	188–211
Solid–liquid equilibria in the ternary system NaBO <sub>2</sub> –NaOH–H <sub>2</sub> O thermal behavior of double salts T. Vilarinho-Franco, A. Teyssier (Grenoble, France and Villeurbanne, France), R. Tenu (Villeurbanne, France), J. Pécaut, J. Delmas, M. Heitzmann, P. Capron (Grenoble, France), J.J. Couniou and C. Goutaudier (Villeurbanne, France).....	212–221
Prediction of the vapor–liquid equilibria and speed of sound in binary systems of 1-alkanols and n-alkanes with the simplified PC-SAFT equation of state X. Liang, K. Thomsen, W. Yan and G.M. Kontogeorgis (Lyngby, Denmark).....	222–232
Liquid–liquid equilibrium data for the system limonene + carvone + ethanol + water at 298.2 K C.M. Oliveira, C.C. Koshima, M.C. Capellini, F.H. Carvalho, K.K. Aracava, C.B. Gonçalves and C.E.C. Rodrigues (Pirassununga, Brazil) .....	233–238
Effect of Eurosemide on denaturation of lysozyme in the presence of ionic surfactant at different temperatures D. Kaushal, D.S. Rana and S. Chauhan (Shimla, India).....	239–247
CaSO <sub>4</sub> solubility in water–ethanol mixtures in the presence of sodium chloride at 25 °C. Application to a reverse osmosis process V. Gomis, M.D. Saquete and J. García-Cano (Alicante, Spain) .....	248–252
Pressure–volume–temperature–composition relations for carbon dioxide + pyrrolidinium-based ionic liquid binary systems T. Makino, M. Kanakubo (Sendai, Japan), T. Umecky (Sendai, Japan and Saga, Japan), A. Suzuki (Sendai, Japan), T. Nishida and J. Takano (Izumiotsu, Japan) .....	253–259
Isobaric vapor–liquid equilibrium at 101.3 kPa and excess properties at 298.15 K for binary mixtures of methyl phenyl carbonate with methanol or dimethyl carbonate I.-C. Hwang, S.-H. Shin, I.-Y. Jeong, Y.-H. Jeon and S.-J. Park (Daejeon, Republic of Korea) .....	260–264
A Raman spectroscopic method for the determination of high pressure vapour liquid equilibria R. Adami (Fisciano, Italy), J. Schuster (Erlangen, Germany), S. Liparoti, E. Reverchon (Fisciano, Italy), A. Leipertz and A. Braeuer (Erlangen, Germany).....	265–273
In situ Raman spectroscopic study of diffusion coefficients of methane in liquid water under high pressure and wide temperatures H. Guo, Y. Chen, W. Lu, L. Li and M. Wang (Wuhan, China) .....	274–278
A group contribution model for determining the vaporization enthalpy of organic compounds at the standard reference temperature of 298 K F. Gharagheizi, P. Ilani-Kashkouli (Durban, South Africa), W.E. Acree Jr. (Denton, TX, USA), A.H. Mohammadi (Durban, South Africa and Paris, France) and D. Ramjugernath (Durban, South Africa).....	279–292
Vapor–liquid equilibria for the 1,1,1,2-tetrafluoroethane (HFC-134a) + 1,1,2,2,3,3-heptafluoropropane (HFC-227ea) and 1,1,1-trifluoroethane (HFC-143a) + 2,3,3,3-tetrafluoroprop-1-ene (HFO-1234yf) systems P. Hu, L.-X. Chen and Z.-S. Chen (Hefei, China) .....	293–297
New correlations between viscosity and surface tension for saturated normal fluids M. Zheng (Qufu, PR China), J. Tian (Qufu, PR China and Dalian, PR China) and Á. Mulero (Badajoz, Spain) .....	298–304
A deduction of the multicriticality conditions of mixtures from the Gibbs tangent plane criterion N. Henderson (Nova Friburgo, Brazil), W.F. Sacco (Santarém, Brazil) and R.A.R. Jr (Nova Friburgo, Brazil) .....	305–308
Spectroscopic and ultrasonic studies on the molecular interaction of certain aromatic amines with p-chloranil R. Kumar, S. Jayakumar and V. Kannappan (Chennai, India) .....	309–319
Equation of state for solid solution–liquid–vapor equilibria at cryogenic conditions S.P. Tan (Tucson, AZ, USA), H. Adidharma (Laramie, WY, USA), J.S. Kargel (Tucson, NV, USA) and G.M. Marion (Reno, NV, USA) ..	320–331
Measurement on the solubility of adipic acid in various solvents at high temperature and its thermodynamics parameters S. Suren, N. Sunsandee (Bangkok, Thailand and Bratislava, Slovak Republic), M. Stolcova, M. Hronec (Bratislava, Slovak Republic), N. Leepipatpiboon, U. Pancharoen and S. Kheawhom (Bangkok, Thailand) .....	332–337
Solubility investigations in the quaternary NaOH–Na <sub>3</sub> VO <sub>4</sub> –Na <sub>2</sub> CrO <sub>4</sub> –H <sub>2</sub> O system at 40 and 80 °C M. Feng, S. Zheng, S. Wang, H. Du and Y. Zhang (Beijing, People's Republic of China) .....	338–342
Determination and correlation of solubilities of 1,2-bis (2-oxo-5,5-dimethyl-1,3,2-dioxyphosphacyclohexyl-2-imino) ethane in selected solvents C.-M. Qi, L.-S. Wang, J.-C. Du, G.-M. Yu and G.-Q. Sun (Beijing, People's Republic of China) .....	343–350
Solute extraction via supercritical ethane from poly(ethylene glycol): A Monte Carlo simulation study J.M. Stubbs (Biddeford, ME, USA) .....	351–356
Salting-out effect of sodium, potassium, carbonate, sulfite, tartrate and thiosulfate ions on aqueous mixtures of acetonitrile or 1-methyl-2-pyrrolidone: A liquid–liquid equilibrium study E. Nemati-Kande (Parsabad, Iran) and H. Shekaari (Tabriz, Iran) .....	357–366

Water + 1-alkanol systems: Modeling the phase, interface and viscosity properties F. Llorell, O. Vilaseca, N. Jung and L.F. Vega (Barcelona, Spain).....	367-378
Liquid-liquid equilibrium of fatty systems: A new approach for adjusting UNIFAC interaction parameters G.F. Hirata (Campinas, Brazil), C.R.A. Abreu (Rio de Janeiro, Brazil), L.C.B.A. Bessa, M.C. Ferreira, E.A.C. Batista and A.J.A. Meirelles (Campinas, Brazil) .....	379-391
Liquid-liquid equilibria studies on ammonium and phosphonium based ionic liquid-aromatic-aliphatic component at $T = 298.15\text{ K}$ and $p = 1\text{ bar}$ : Correlations and a-priori predictions C.V. Manohar, D. Rabari, A.A.P. Kumar, T. Banerjee and K. Mohanty (Guwahati, India).....	392-400
Improved hydrodynamic equations for the accurate prediction of diffusivities in supercritical carbon dioxide R.V. Vaz, A.L. Magalhães and C.M. Silva (Aveiro, Portugal).....	401-415
Phase equilibria of binary mixtures (ionic liquid + aromatic hydrocarbon): Effect of the structure of the components on the solubility E.J. González (Porto, Portugal), P.F. Requejo, Á. Domínguez (Vigo, Spain) and E.A. Macedo (Porto, Portugal).....	416-422
Molecular modeling of the ideal gas enthalpy of formation of hydrocarbons T.N.G. Borhani (Skudai, Malaysia), M. Bagheri (Tehran, Iran) and Z.A. Manan (Skudai, Malaysia) .....	423-434
Behavior of Triton X-114 cloud point in the presence of inorganic electrolytes V.C. Santos-Ebinuma (Araraquara, Brazil and São Paulo, Brazil), A.M. Lopes (São Paulo, Brazil), A. Converti (Genoa, Italy), A.P. Júnior and C. de Oliveira Rangel-Yagui (São Paulo, Brazil) .....	435-438
Vapor-liquid equilibrium for tetrahydrofuran + methanol + tetrafluoroborate-based ionic liquids at 101.3 kPa Q. Li, P. Liu, L. Cao, F. Wen, S. Zhang and B. Wang (Beijing, China).....	439-444
Experimental data on xanthohumol solubility in supercritical carbon dioxide D. Kostrzewska, A. Dobrzańska-Inger and E. Rój (Puławy, Poland) .....	445-450
Isobaric vapor-liquid equilibrium for binary systems of <i>n</i> -undecane + p-, o-, m-xylene at 10 kPa S. Yang, F. Zhu, Y. Zhu and M. Qiao (Xinxiang, PR China) .....	451-455
Methane and carbon dioxide adsorption in clay-like slit pores by Monte Carlo simulations Z. Jin and A. Firoozabadi (USA).....	456-465
Measurement and correlation solubility and mixing properties of L-malic acid in pure and mixed organic solvents Y. Kai, Y. Hu, Z. Cao, X. Liu, Y. Liu and W. Yang (Nanjing, China).....	466-471
Gibbs ensemble simulation on polarizable models: Vapor-liquid equilibrium in Baranyai-Kiss models of water F. Moučka (Ústí nad Labem, Czech Republic) and I. Nezbeda (Ústí nad Labem, Czech Republic and Prague 6, Czech Republic)...	472-476