


M  
F70



Volume 362, January 25 2014

ISSN 0378-3812

Special Issue: PPEPPD 2013

Guest Editor: Susana Bottini

**FLUID PHASE**

**EQUILIBRIA**

**AN INTERNATIONAL JOURNAL**

**FLUID PHASE**

**EQUILIBRIA**

## CONTENTS

(Abstracted/Indexed in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®*)

Novel aqueous two-phase system based on a hyperbranched polymer A. Kulaguin-Chicaroux and T. Zeiner (Dortmund, Germany) .....	1-10
Solubility of solids in supercritical fluid using the hard-body expanded virial equation of state S.J. Pai and Y.C. Bae (Seoul, Republic of Korea) .....	11-18
Measuring and correlating diffusivity in polymer-solvent systems using free-volume theory R.P. Danner (University Park, PA, USA) .....	19-27
Experimental and modeling investigation of kinetics of methane gas hydrate formation in water-in-oil emulsion L. Mu, S. Li, Q.-L. Ma, K. Zhang, C.-Y. Sun, G.-J. Chen, B. Liu and L.-Y. Yang (Beijing, China) .....	28-34
Correlation of phase equilibria for the systems containing 1-butanol + water by concentration dependent surface area parameter model Y. Iwai and I. Taniguchi (Fukuoka, Japan) .....	35-40
PC-SAFT parameters from ab initio calculations M. Umer (Aachen, Germany), K. Albers, G. Sadowski (Dortmund, Germany) and K. Leonhard (Aachen, Germany) .....	41-50
Robust and efficient Trust-Region based stability analysis and multiphase flash calculations M. Petitfrere and D.V. Nichita (Pau Cedex, France) .....	51-68
Study on behavior of alkaline fiber FFA-1 adsorbing glyphosate from production wastewater of glyphosate C.R. Zhou, G.P. Li and D.G. Jiang (Zhengzhou, PR China) .....	69-73
Prediction of physical properties for molecular design of solvents J. Scilipoti, M. Cismondi (Córdoba, Argentina) and E.A. Brignole (Bahía Blanca, Argentina) .....	74-80
Hydrophobicity and geometry: Water at curved graphitic-like surfaces and within model pores in self-assembled monolayers L.M. Alarcón, J.M. Montes de Oca, S.R. Accordino, J.A. Rodriguez Fris and G.A. Appignanesi (Bahía Blanca, Argentina) .....	81-86
Prediction of solid polycyclic aromatic hydrocarbons solubility in water, with the NRTL-PR model J. Escandell, I. Raspo and E. Neau (Marseille, France) .....	87-95
High pressure phase equilibria of binary mixtures of light hydrocarbons in the ionic liquid 1-hexyl-3-methylimidazolium tetracyanoborate M.T. Mota-Martinez, M. Althuluth (Abu Dhabi, United Arab Emirates and Eindhoven, The Netherlands), A. Berrouk (Abu Dhabi, United Arab Emirates), M.C. Kroon (Eindhoven, The Netherlands and C.J. Peters (Abu Dhabi, United Arab Emirates and Eindhoven, The Netherlands) .....	96-101
Some examples of the contribution of applied thermodynamics to Post-Combustion CO <sub>2</sub> -Capture technology P.M. Mathias (Aliso Viejo, CA, USA) .....	102-107
Thermodynamic model for process design, simulation and optimization in the production of biodiesel N. Cotabarren, P. Hegel and S. Pereda (Bahía Blanca, Argentina) .....	108-112
Influences on the stability of collagen triple-helix S. Schweizer, A. Bick, L. Subramanian and X. Krokidis (Paris, France) .....	113-117
Concentration effects on the selective extraction of ethanol from aqueous solution using silicalite-1 and decanol isomers S.J. Keasler, P. Bai, M. Tsapatsis and J.I. Siepmann (Minneapolis, MN, USA) .....	118-124
High pressure phase behavior modeling of asymmetric alkane + alkane binary systems with the RKPR EOS M. Cismondi Duarte (Argentina and Córdoba, Argentina), M.V. Galdo, M.J. Gomez, N.G. Tassin and M. Yanes (Córdoba, Argentina) .....	125-135
Evaluation of different contribution methods over the performance of Peng-Robinson and CPA equation of state in the correlation of VLE of triglycerides, fatty esters and glycerol + CO <sub>2</sub> and alcohol S. Arvelos, L.L. Rade, E.O. Watanabe, C.E. Hori and L.L. Romanielo (Uberlândia, Brazil) .....	136-146
Experimental phase behavior study of a five-component model gas condensate A. Shariati (Shiraz, Iran), E.J.M. Straver, L.J. Florusse (Delft, The Netherlands) and C.J. Peters (Abu Dhabi, United Arab Emirates, Eindhoven, The Netherlands) .....	147-150
Interfacial tension of binary mixtures exhibiting azeotropic behavior: Measurement and modeling with PCP-SAFT combined with Density Gradient Theory E. Schäfer (Berlin, Germany and Dortmund, Germany), G. Sadowski (Dortmund, Germany) and S. Enders (Berlin, Germany) .....	151-162

(Contents continued on BM I)



(contents continued)

Liquid-liquid equilibria of binary systems {benzene + [x-Mim][NTf <sub>2</sub> ] ionic liquid}: Experimental data and thermodynamic modeling using a group contribution equation of state E.J. González (Porto, Portugal), S.B. Bottini, S. Pereda (Bahía Blanca, Argentina) and Eugénia.A. Macedo (Porto, Portugal).....	163-169
Phase behavior and mesoscale solubilization in aqueous solutions of hydrotropes D. Subramanian and M.A. Anisimov (College Park, MD, USA).....	170-176
Steric effects on ion dynamics near charged electrodes P.H.R. Alijó, F.W. Tavares, E.C. Biscaia Jr. and A.R. Secchi (Rio de Janeiro, Brazil).....	177-186
Effect of solvation with salting effect on solubilities of fluorocarbons in alcohols H. Nishiumi (Koganei-city, Japan), K.-i. Ago (Chiyoda-ku, Japan) and D. Kodama (Koriyamashi, Japan).....	187-191
From conventional to more sustainable fuels Trends and needs in research in the thermodynamics area S. Jullian and X. Longaygue (Rueil-Malmaison, France).....	192-195
New formulation of the lattice cluster theory equation of state for multi-component systems K. Langenbach (Berlin, Germany), D. Browarzik (Halle, Germany), J. Sailer and S. Enders (Berlin, Germany).....	196-212
Generation of characteristic maps of the fluid phase behavior of ternary systems G. Pisoni (Bahía Blanca, Argentina), M. Cismondi (Argentina and Córdoba, Argentina), L. Cardozo-Filho (Maringá, Brazil) and M.S. Zabaloy (Bahía Blanca, Argentina).....	213-226
Adsorption and separation of CO <sub>2</sub> /CH <sub>4</sub> mixtures using nanoporous adsorbents by molecular simulation L. Lu, S. Wang (Nanjing, PR China), E.A. Müller (London, UK), W. Cao, Y. Zhu, X. Lu (Nanjing, PR China) and G. Jackson (London, UK).....	227-234
Flow resistance analysis of nanoconfined water in silt pores by molecular simulations: Effect of pore wall interfacial properties Y. Zhu (Nanjing, PR China), L. Zhang (Champaign, IL, USA), X. Lu, L. Lu and X. Wu (Nanjing, PR China).....	235-241
Accurately modeling benzene and alkylbenzenes using a group contribution based SAFT approach G. Das, M.C.d. Ramos and C. M <sup>c</sup> Cabe (Nashville, TN, USA).....	242-251
Hydrogen separation via forming hydrate in W/O emulsion B. Liu, H. Liu, B. Wang, J. Wang, C. Sun, X. Zeng and G. Chen (Beijing, PR China).....	252-257
An equation of state for solid-liquid-vapor equilibrium applied to gas processing and natural gas liquefaction P. Stringari, M. Campestrini, C. Coquelet (Fontainebleau, France) and P. Arpentinier (Jouy-en-Josas, France).....	258-267
Thermodynamic properties of aqueous solutions of single and multiple salts using the Q-electrolattice equation of state A. Zuber (Doha, Qatar and Paraná, Brazil), R.F. Checoni and M. Castier (Doha, Qatar).....	268-280
Salt-forming regions of seawater type solution in the evaporation and fractional crystallization process H. Zhou, Y. Bao, X. Bai, R. Ma, L. Huangfu and C. Zhang (Tianjin, PR China).....	281-287
A comprehensive framework for surfactant selection and design for emulsion based chemical product design M. Mattei, G.M. Kontogeorgis and R. Gani (Lyngby, Denmark).....	288-299
CO <sub>2</sub> absorption properties, densities, viscosities, and electrical conductivities of ethylimidazolium and 1-ethyl-3-methylimidazolium ionic liquids T. Makino, M. Kanakubo, Y. Masuda (Sendai, Japan), T. Umecky (Saga, Japan) and A. Suzuki (Sendai, Japan).....	300-306
Liquid-vapor equilibrium data of CO <sub>2</sub> + dichloromethane + medroxyprogesterone system W.M. Giufrida, L.F. Pinto, A.F. Zanette (Maringá, Brazil), F.A.P. Voll (Curitiba, Brazil), M.H. Kunita, V.F. Cabral and L. Cardozo-Filho (Maringá, Brazil).....	307-312
Determination and correlation of excess molar enthalpies of the binary carbon dioxide + ethyl acetate system at 298.15 to 308.15 K and 5.0 to 7.0 MPa K. Kurihara, Y. Nakamura, T. Okamoto, Y. Itazu, H. Matsuda and K. Tochigi (Tokyo, Japan).....	313-317
Data, analysis and modeling of physical properties for process design of systems involving lipids L.P. Cunico (Kongens Lyngby, Denmark), R. Ceriani (Campinas, Brazil), B. Sarup (Soeborg, Denmark), J.P. O'Connell (Charlottesville, VA, USA) and R. Gani (Kongens Lyngby, Denmark).....	318-327
Towards automated characterisation of liquid-liquid equilibria D. Dechambre, C. Pauls (Aachen, Germany), L. Greiner (Mannheim, Germany), K. Leonhard and A. Bardow (Aachen, Germany)...	328-334
Phase equilibrium of fluids confined in porous media from an extended Peng-Robinson equation of state L. Travalloni (Rio de Janeiro, Brazil), M. Castier (Doha, Qatar) and F.W. Tavares (Rio de Janeiro, Brazil).....	335-341
Designing new amine functionalized metal-organic frameworks for carbon dioxide/methane separation Z. Qiao, J. Zhou (Guangzhou, PR China) and X. Lu (Nanjing, PR China).....	342-348
Molecular simulations of myoglobin adsorbed on rutile (1 1 0) and (0 0 1) surfaces C. Yang, C. Peng, D. Zhao, C. Liao, J. Zhou (Guangzhou, PR China) and X. Lu (Nanjing, PR China).....	349-354
New method to detect asphaltene precipitation onset induced by CO <sub>2</sub> injection N.A.E. da Silva, V.R.d.R. Oliveira, M.M.S. Souza, Y. Guerrieri and G.M.N. Costa (Salvador, Brazil).....	355-364
Multiphase equilibrium calculations using the semi-continuous thermodynamics of hydrocarbon mixtures M. Petitfrere, D.V. Nichita and F. Montel (Pau Cedex, France).....	365-378
Phase equilibrium relations for binary mixed hydrate systems composed of carbon dioxide and cyclopentane derivatives Y. Matsumoto (Osaka, Japan), T. Makino (Miyagi, Japan), T. Sugahara and K. Ohgaki (Osaka, Japan).....	379-382
Hydrogen solubility in triolein, and propane solubility in oleic acid for second generation BDF synthesis by use of hydrodeoxygenation reaction T. Tsuji, K.-h. Ohya, T.-a. Hoshina, T. Hiaki (Narashino, Japan), K. Maeda (Himeji, Japan), H. Kuramochi and M. Osako (Tsukuba, Japan).....	383-388