


ПН

F70



Volume 363, February 15 2014

ISSN 0378-3812

**FLUID PHASE**

**EQUILIBRIA**

**AN INTERNATIONAL JOURNAL**

**FLUID PHASE**

**EQUILIBRIA**

## CONTENTS

(Abstracted/Indexed in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®*)

**Full Length Article**

Prediction of true critical temperature of multi-component mixtures: An extension to Chueh and Prausnitz method H. Najafi, B. Maghbooli and M.A. Sobati (Tehran, Iran).....	1–17
Development and evaluation of a new semi-empirical model for correlation of drug solubility in supercritical CO <sub>2</sub> K. Keshmiri, A. Vatanara and Y. Yamini (Tehran, Iran).....	18–26
Are the reservoir fluid compositional grading data reliable? A.H. Mohammadi (Paris, France and Durban, South Africa), A. Eslamimanesh (Potsdam, NY, USA), F. Gharagheizi and P. Ilani-Kashkouli (Durban, South Africa and Buinzahra, Iran).....	27–31
Study of thermodynamic properties of L-serine in aqueous L-carboxymethyl-3-methylimidazolium chloride solutions at 298.15 K M.T. Zafarani-Moattar, B. Asadzadeh, A. Shahrissa and M.G. Nazari (Tabriz, Iran).....	32–40
Extraction of toluene from alkane using [Bmim][NO <sub>3</sub> ] or [Omim][NO <sub>3</sub> ] ionic liquid at 298.15 K and atmospheric pressure B. Mokhtarani (Tehran, Iran), J. Musavi and M. Parvini (Semnan, Iran).....	41–47
Free volume investigation of imidazolium ionic liquids from positron lifetime spectroscopy Y. Yu (Nanjing, China and Halle, Germany), D. Bejan (Wuppertal, Germany) and R. Krause-Rehberg (Halle, Germany) ...	48–54
Solid-liquid equilibrium of quaternary system Cd <sup>2+</sup> , K <sup>+</sup> , Na <sup>+</sup> //SO <sub>4</sub> <sup>2-</sup> -H <sub>2</sub> O at 298 K W. Zhang, Y. Huang, Y. Chen (Chengdu, PR China), S.S. Hughes (USA), L.-c. Zhong and F. Zou (Chengdu, PR China).....	55–58
Modeling imidazolium-based ionic liquids with ePC-SAFT. Part II. Application to H <sub>2</sub> S and synthesis-gas components X. Ji (Luleå, Sweden), C. Held and G. Sadowski (Dortmund, Germany).....	59–65
The fluidity of room temperature ionic liquids Y. Marcus (Jerusalem, Israel).....	66–69
Experimental measurement and prediction of dissociation condition for carbon dioxide clathrate hydrates in the presence of methanol/ethylene glycol + sodium chloride/magnesium chloride aqueous solution H. Najibi, E. Amiri (Ahwaz, Iran) and A.H. Mohammadi (Paris, France).....	70–73
Performance comparison of adsorption isotherm models for supercritical hydrogen sorption on MOFs E. Dundar, R. Zacharia, R. Chahine and P. Bénard (Trois-Rivières, Canada).....	74–85
Solubility of androstenedione in lower alcohols W. Tang, C. Xie, Z. Wang, S. Wu, Y. Feng, X. Wang, J. Wang and J. Gong (Tianjin, China).....	86–96
Solubility of carbamazepine, nicotinamide and carbamazepine-nicotinamide cocrystal in ethanol-water mixtures A. Shayanfar (Tabriz, Iran), S. Velaga (Lulea, Sweden) and A. Jouyban (Tabriz, Iran).....	97–105
Thermodynamic and aggregation properties of sodium n-hexylsulfonate in aqueous solution S. Solaimani and R. Sadeghi (Sanandaj, Iran).....	106–116
Measurement and correlation of the mutual solubility of diisopropylamine + water and triethylamine + water systems at high pressure Y. Tian, P. Yuan, L. Cao and R. Zhu (Tianjin, China).....	117–120
Application of constrained multi-variable search methods for prediction of PVT properties of crude oil systems M. Arabloo, M.-A. Amooie, A. Hemmati-Sarapardeh, M.-H. Ghazanfari (Tehran, Iran) and A.H. Mohammadi (Paris, France and Durban, South Africa).....	121–130
( <i>p</i> , <i>V</i> <sup>o</sup> , <i>T</i> ) Measurements of mixtures (DBE + alcohol) at temperatures from (293.15 to 353.15) K and at pressures up to 140 MPa F.E.M. Alaoui (Pau, France, Burgos, Spain and El Jadida Plateau, Morocco), E.A. Montero (Burgos, Spain), J.-P. Bazile (Pau, France), F. Aguilar (Burgos, Spain) and C. Boned (Pau, France).....	131–148
Effect of impurities in captured CO <sub>2</sub> on liquid-vapor equilibrium M. Ahmad (Groningen, The Netherlands), J. Gernert (Bochum, Germany) and E. Wilbers (Groningen, The Netherlands)....	149–155
Excess molar volumes and excess molar enthalpies in binary systems N-alkyl-triethylammonium bis(trifluoromethylsulfonyl) imide + methanol K. Machanová (Prague, Czech Republic), J. Troncoso (Ourense, Spain), J. Jacquemin (Belfast, UK) and M. Bendová (Prague, Czech Republic).....	156–166

(Contents continued on BM I)



(contents continued)

Experimentation and thermodynamic representations of binaries containing compounds of low boiling points: Pentane and alkyl methanoates L. Fernández, J. Ortega, G. Sabater and F. Espiau (Las Palmas de Gran Canaria, Spain) .....	167-179
Mutual diffusion coefficients, density, and viscosity of aqueous solutions of new polyamine CO <sub>2</sub> absorbents C.-P. Liao (Chung Li, Taiwan), R.B. Leron (Chung Li, Taiwan and Manila, Philippines) and M.-H. Li (Chung Li, Taiwan) .....	180-188
Development of a new generalized correlation to characterize physical properties of pure components and petroleum fractions P. Hosseinfar and S. Jamshidi (Tehran, Iran) .....	189-198
Liquid-liquid equilibrium of the CuSO <sub>4</sub> + PEG 4000 + H <sub>2</sub> O system at different temperatures M. Claros, M.E. Taboada, H.R. Galleguillos and Y.P. Jimenez (Antofagasta, Chile) .....	199-206
A modified Hoch-Arps Hofen model for the calculation of enthalpies of formation of multicomponent liquid alloys X. Yang, H. Yang, D. Tao and D. Liu (Kunming, PR China) .....	207-212
Prediction of volumetric properties of polymer melts using new perturbed hard-chain equation of state S.M. Hosseini, M.M. Alavianmehr and J. Moghadasi (Shiraz, Iran) .....	213-219
Isobaric vapor-liquid equilibrium for water + acetic acid + 1-butyl-3- methylimidazolium dibutylphosphate at 101.32 kPa Y. Peng, J. Fu and X. Lu (Hangzhou, PR China) .....	220-227
Solid-liquid equilibrium of dicyandiamide in different solvents X. Zhang, G. Qian, X. Yang, C. Hu and X. Zhou (Shanghai, China) .....	228-232
A priori prediction of the octanol-water partition coefficient ( $K_{ow}$ ) of ionic liquids B.-S. Lee and S.-T. Lin (Taiwan, Taiwan) .....	233-238
Extended hard-sphere model for the viscosity of dense fluids F. Ciotta, J.P.M. Trusler and V. Vesovic (London, UK) .....	239-247
Prediction and measurement of phase equilibria for the extraction of BTX from naphtha reformat using BMIMPF <sub>6</sub> ionic liquid O.A. Al-Rashed, M.A. Fahim and M. Shaaban (Safat, Kuwait) .....	248-262
High pressure measurement and thermodynamic modeling the solubility of H <sub>2</sub> S in the aqueous <i>N</i> -methyldiethanolamine + 2-amino-2-methyl-1-propanol + piperazine systems A. Haghtalab, A. Izadi and A. Shojaeian (Tehran, Iran) .....	263-275
Solubility and solution thermodynamics of 2,3,4,5-tetrabromothiophene in (ethanol + trichloromethane) binary solvent mixtures W. Yang, C. Wang, Y. Hu, Y. Li and K. Wang (Nanjing, China) .....	276-281
Temperature dependence and the effect of charge configuration on water permeation through modified carbon nanotubes: A simulation study A. Alizadeh and G. Parsafar (Tehran, Iran) .....	282-289
<b>Short Communication</b>	
On solving the Rachford-Rice equation with higher order methods W. Yan and E.H. Stenby (Kgs. Lyngby, Denmark) .....	290-292