

NU
F70



Volume 369, May 15 2014

ISSN 0378-3812

FLUID PHASE EQUILIBRIA

AN INTERNATIONAL JOURNAL

FLUID PHASE EQUILIBRIA

CONTENTS

(Abstracted/Index in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr.*, ASCA, Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®)

Full Length Articles

Prediction of hydrate formation conditions using G ^E -EOS and UNIQUAC models for pure and mixed-gas systems H. Delavar and A. Haghtalab (Tehran, Iran)	1–12
Helmholtz energy and extended corresponding states model for the prediction of thermodynamic properties of refrigerants J.F. Estela-Uribe (Cali, Colombia)	13–32
Phase equilibrium properties of binary and ternary mixtures containing 2-butanol, 2,2,4-trimethylpentane and 1-hexene at 313.15 K A. Belabbaci, S. Ghellai (Tlemcen, Algeria), R.M. Villamañán, M.C. Martín (Valladolid, Spain), L. Negadi (Tlemcen, Algeria) and M.A. Villamañán (Valladolid, Spain)	33–38
Rheological behavior and physico-chemical properties of cetyltrimethylammonium salicylate in aqueous solutions A. Ping, P. Du, X. Wei, D. Sun, J. Zhang, J. Liu and X. Zhong (Liaocheng, PR China)	39–46
Phase equilibria of methane clathrate hydrates from Grand Canonical Monte Carlo simulations M. Lasich (Durban, South Africa), A.H. Mohammadi (Durban, South Africa and Paris, France), K. Bolton (Borås, Sweden), J. Vrabec (Paderborn, Germany) and D. Ramjugernath (Durban, South Africa)	47–54
Evaluation of COSMO-RS model in binary and ternary mixtures of natural antioxidants, ionic liquids and organic solvents E.I. Alevizou and E.C. Voutsas (Athens, Greece)	55–67
Thermodynamic properties of binary aqueous solutions of orthophosphate salts, sodium, potassium and ammonium at T = 298.15 K M. El Guendouzi and A. Benbiyi (Morocco)	68–85
PC-SAFT modeling of asphaltene phase behavior in the presence of nonionic dispersants M. Sedghi and L. Goual (Laramie, WY, USA)	86–94
VLE and LLE in ternary systems of two associating components (water, aniline, and cyclohexylamine) and a hydrocarbon (cyclohexane or methylcyclohexane) M. Klauck, R. Silbermann, R. Metasch, T. Jasinski, G. Kalies and J. Schmelzer (Dresden, Germany)	95–108
Isobaric (vapour + liquid) equilibria for three binary systems (toluene + anisole, n-butylbenzene + anisole, and guaiacol + anisole) at 101.33 kPa H. Li, S. Xia, P. Ma, F. Yan, Z. Yang and Y. Li (Tianjin, People's Republic of China)	109–114
Letter to the Editor	
Comment on "The solubility of mercaptosuccinic acid in water + (methanol, ethanol, acetone) mixtures from (278.15 to 333.15 K). <i>Fluid Phase Equilib.</i> 361 (2014) 282–288." J.-Q. Liu (Luoyang, PR China)	115–118

