



Volume 375, August 15 2014

ISSN 0378-3812

FLUID PHASE EQUILIBRIA

AN INTERNATIONAL JOURNAL

FLUID PHASE EQUILIBRIA

CONTENTS

(Abstracted/Indexed in: Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®)

Full Length Articles

- Determination and thermodynamic evaluation of isobaric VLE of methyl acetate or ethyl acetate with 2-propanol at 0.3 and 0.6 MPa
P. Susial, R. Susial, E.J. Estupiñan, V.D. Castillo, J.J. Rodríguez-Henríquez and J.C. Apolinario (Las Palmas de Gran Canaria, Spain) 1–10
- Theory-based constraints on variations of infinite dilution partial molar volumes of aqueous solutes at various temperatures and water densities
A.V. Plyasunov (Moscow Region, Russia) 11–17
- Partial molar volumes and compressibilities of glycine betaine in aqueous NaCl solutions at temperatures $T = (288.15\text{--}318.15)$ K
V.R. Karanth and D.K. Bhat (Surathkal, India) 18–22
- Modulating the aggregation behavior of aqueous sodium dodecylsulphate (SDS) with addition of trisubstituted imidazolium based ionic liquid 1-butyl-2,3-dimethylimidazolium tetrafluoroborate [bdmim][BF₄]
A. Pal and A. Pillania (Kurukshetra, India) 23–29
- Effect of ionic liquids as adjuvants on PEG-based ABS formation and the extraction of two probe dyes
R.L. de Souza, V.C. Campos (Aracaju-SE, Brazil), S.P.M. Ventura (Aveiro, Portugal), C.M.F. Soares (Aracaju-SE, Brazil), J.A.P. Coutinho (Aveiro, Portugal) and Á.S. Lima (Aracaju-SE, Brazil) 30–36
- Isobaric vapor–liquid equilibrium for binary and ternary systems with toluene, 2-methoxyethanol and dimethyl sulfoxide at 101.3 kPa
S. Zhao, P. Bai and C. Sun (Tianjin, PR China) 37–44
- Modelling of tetrahydrofuran promoted gas hydrate systems for carbon dioxide capture processes
P.J. Herslund, K. Thomsen, J. Abildskov and N. von Solms (Kgs. Lyngby., Denmark) 45–65
- Solubility, liquid–liquid equilibrium and critical states for the system acetic acid + *n*-propanol + *n*-propyl acetate + water at 293.15 K and 303.15 K
M. Toikka, A. Samarov and A. Toikka (St. Petersburg, Russia) 66–72
- Thermodynamics of risperidone and solubility in pure organic solvents
D. Mealey (Castletroy, Ireland), M. Svärd and Å.C. Rasmuson (Castletroy, Ireland and Stockholm, Sweden) 73–79
- The effect of the cation aromaticity upon the thermophysical properties of piperidinium- and pyridinium-based ionic liquids
A. Bhattacharjee, P.J. Carvalho and J.A.P. Coutinho (Aveiro, Portugal) 80–88
- Modelling of cyclopentane promoted gas hydrate systems for carbon dioxide capture processes
P.J. Herslund, K. Thomsen, J. Abildskov and N. von Solms (Lyngby., Denmark) 89–103
- Stability of clavulanic acid in PEG/citrate and liquid–liquid extraction in aqueous two-phase system
M.N. Carneiro-da-Cunha, K.P.S. Souza (Recife-PE, Brazil), A.M.O. Mota, J.A. Teixeira (Braga, Portugal), C.S. Porto (Recife-PE, Brazil), T.S. Porto (Recife-PE, Brazil and Garanhuns-PE, Brazil) and A.L.F. Porto (Recife-PE, Brazil) 104–109
- Solid–liquid equilibrium and thermodynamic of 2,5-thiophenedicarboxylic acid in different organic solvents
W. Zhi, Y. Hu, M. Liang, Y. Liu, J. Li, J. Yin (Nanjing, China) and Y. Shi (Taiyuan, China) 110–114
- Solubility of benzyl methacrylate/methacrylic acid copolymers in carbon dioxide-expanded acetone and correlation with the PC-SAFT equation of state
S.M. Paap, K. Frey, G.R. Johnson and J.W. Tester (Cambridge, MA, USA) 115–123
- Excess enthalpies for pseudobinary mixtures containing vegetable oils at the temperatures 298.15 K, 353.15 K and 383.15 K
P.C. Belting (Oldenburg, Federal Republic of Germany and Campinas-SP, Brazil), J. Gmehling, R. Bölts, J. Rarey (Oldenburg, Federal Republic of Germany), R. Ceriani (Campinas-SP, Brazil), O. Chiavone-Filho (Natal-RN, Brazil) and A.J.A. Meirelles (Campinas-SP, Brazil) 124–133
- Solubility of CO₂/CH₄ gas mixtures in ionic liquids
M. Ramdin, A. Amlianitis, T.W. de Loos and T.J.H. Vlucht (Delft, The Netherlands) 134–142
- Prediction of isenthalps, Joule–Thomson Coefficients and Joule–Thomson inversion curves of refrigerants by molecular simulation
S. Figueroa-Gerstenmaier (Leon, Mexico), M. Lísal (Ústí n. Lab., Czech Republic and Prague 6, Czech Republic), I. Nezbeda (Prague 6, Czech Republic and Ústí n. Lab., Czech Republic), W.R. Smith (Oshawa, Canada and Guelph, Canada) and V.M. Trejos (Leon, Mexico) 143–151

(Contents continued on BM I)



(contents continued)

Testing solubility data of H ₂ S and SO ₂ in ionic liquids for sulfur-removal processes C.A. Faúndez, J.F. Díaz-Valdés (Concepción, Chile) and J.O. Valderrama (La Serena, Chile)	152-160
Impact of the cation symmetry on the mutual solubilities between water and imidazolium-based ionic liquids M.A.R. Martins, C.M.S.S. Neves, K.A. Kurnia, A. Luís (Aveiro, Portugal), L.M.N.B.F. Santos (Porto, Portugal), M.G. Freire (Aveiro, Portugal), S.P. Pinho (Bragança, Portugal and Salvador, Brazil) and J.A.P. Coutinho (Aveiro, Portugal)	161-167
Electrolyte-cosolvent effects on the properties of micellar and monolayer phases in the cationic-rich region of catanionic mixture: The phase transition between microstructures and nanostructures B. Sohrabi, M. Moallemi, R. Amani and M. Kiasadegh (Tehran, Iran)	168-175
High pressure vapor-liquid equilibria measurements and modeling of butane/ethanol system and isobutane/ethanol system H. Machida, K. Matsumura and H. Horizoe (Nagoya, Japan)	176-180
Simultaneous measurement solubility of carbon dioxide + hydrogen sulfide into aqueous blends of alkanolamines at high pressure A. Haghtalab and A. Izadi (Tehran, Iran)	181-190
Toward mechanistic understanding of heavy crude oil/brine interfacial tension: The roles of salinity, temperature and pressure F. Moeini, A. Hemmati-Sarapardeh, M.-H. Ghazanfari, M. Masihi and S. Ayatollahi (Tehran, Iran)	191-200
Vapour-liquid equilibrium of propionic acid + caproic acid, isobutyric acid + caproic acid, valeric acid + caproic acid and caproic acid + enanthoic acid binary mixtures A. Hlengwere, S.A. Iwarere, P. Naidoo, J.D. Raal and D. Ramjugernath (Durban, South Africa)	201-208
Calculation of phase equilibria for multi-component mixtures using highly accurate Helmholtz energy equations of state J. Gernert, A. Jäger and R. Span (Bochum, Germany)	209-218
Density and viscometric study of binary liquid mixtures of morpholine with some aromatic hydrocarbons at temperatures 303.15, 308.15 and 313.15 K S. Sharma and M. Makavana (Patan, India)	219-227
Thermal conductivity of ethyl fluoride (HFC161) C. Yao, X. Zhao, S. Lv (Xi'an, China) and Z. Guo (Zhejiang, China)	228-235
A new model based on multilayer kinetic adsorption mechanism for asphaltene adsorption in porous media during dynamic condition T. Jafari Behbahani, C. Ghorbi, V. Taghikhani and A. Shahrabadi (Tehran, Iran)	236-245
An experimental investigation on the solubility of CO ₂ in phenol M. Jödecke, Á. Pérez-Salado Kamps, J. Xia and G. Maurer (Kaiserslautern, Germany)	246-253
On petroleum fluid characterization with the PC-SAFT equation of state X. Liang, W. Yan, K. Thomsen and G.M. Kontogeorgis (Lyngby, Denmark)	254-268
The hydrogen bonding analysis of 1-methylimidazole-water mixtures studied by dielectric relaxation spectroscopy G.-Z. Jia, J.-C. Liu and X.-Q. Yang (Chengdu, China)	269-274
Flash-point estimation for binary partially miscible mixtures of flammable solvents by UNIFAC group contribution methods H.-J. Liaw and T.-P. Tsai (Kaohsiung, Taiwan)	275-285
A conductance study to analyze the effect of organic solvents on micellization behavior of carbohydrate-surfactant system at variable temperatures S. Chauhan, J. Jyoti, K. Sharma and K. Kumar (Shimla, India)	286-292
Prediction of VOCs adsorption equilibria on activated carbon in supercritical carbon dioxide over a wide range of temperature and pressure by using pure component adsorption data: Combined approach of the Dubinin-Astakhov equation and the non-ideal adsorbed solution theory (NAIST) I. Ushiki, M. Ota, Y. Sato and H. Inomata (Sendai, Japan)	293-305
Modeling aqueous two-phase systems: II. Inorganic salts and polyether homo- and copolymers as ATPS former T. Reschke, C. Brandenbusch and G. Sadowski (Dortmund, Germany)	306-315
Series approach to modeling of amino acid and electrolyte mixtures in the aqueous media M. Alamdari Shendi (Tabriz, Iran), Z. Karimzadeh (Evin-Tehran, Iran) and R. Salamat-Ahangari (Tabriz, Iran)	316-323
Effect of cation on the solubility of carbon dioxide in three bis(fluorosulfonyl)imide low viscosity ([FSI]) ionic liquids A. Tagiuri, K.Z. Sumon, A. Henni (Regina, Canada), K. Zanganeh and A. Shafeen (Ottawa, Canada)	324-331
A simple correlation to predict drug solubility in supercritical carbon dioxide A.A. Amooey (Babolsar, Iran)	332-339
Solubilization efficiency of mixed cationic aggregates towards aromatic compounds A. Kumar, G. Kaur, G.R. Chaudhary and S.K. Mehta (Chandigarh, India)	340-346
Temperature dependence of limiting activity coefficients, Henry's law constants, and related infinite dilution properties of C ₄ -C ₆ isomeric <i>n</i> -alkyl ethanoates/ethyl <i>n</i> -alkanoates in water. Measurement, critical compilation, correlation, and recommended data D. Fenclová, A. Blahut, P. Vrbka, V. Dohnal (Prague 6, Czech Republic) and A. Böhme (Leipzig, Germany)	347-359
A note on effective phase stability calculations using a Gradient-Based Cuckoo Search algorithm S.K. Fateen (Egypt) and A. Bonilla-Petriciolet (Mexico)	360-366
The surface chemical potential from a surface equation of state versus Butler's equation C. Bermúdez-Salguero and J. Gracia-Fadrique (Mexico)	367-372
Corrigendum to "The solubility of mercaptosuccinic acid in water + (methanol, ethanol, acetone) mixtures from (278.15 to 333.15 K)" [Fluid Phase Equilib. 361 (2014) 282-288] Y. Kai, Y. Hu, Y. Liu, M. Liang, W. Yang and Y. Xi (Nanjing, China)	373-374