


FU
E70

Volume 379, October 15 2014

ISSN 0378-3812



FLUID PHASE EQUILIBRIA

AN INTERNATIONAL JOURNAL

FLUID PHASE EQUILIBRIA

CONTENTS

(Abstracted/Indexed in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA, Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®*)

Full length Articles

Miscibility of ternary systems containing kerosene-based surrogate fuel and hydrous ethanol: Experimental data + thermodynamic modeling L. Chen, P. Sun, S. Ding and S. Yang (China)	1-9
Solubility and tie-line data for ternary aqueous mixtures of cyclopentanol with organic solvents at $T = 298.2$ K: Experiments and NRTL model T. Wongsawa (Bangkok, Thailand), M. Hronec (Bratislava, Slovakia), U. Pancharoen and S. Phatanasri (Bangkok, Thailand) ...	10-18
Liquid-liquid equilibria of water + lactic acid + methyl isobutyl ketone T.J. Afolabi and A.I. Alao (Ogbomoso, Nigeria)	19-25
A robust algorithm for isenthalpic flash of narrow-boiling fluids D. Zhu and R. Okuno (Canada)	26-51
Isothermal vapor-liquid equilibria for the binary system of dimethyl ether (DME) + methyl iodide (CH_3I) H. Cho, J.E. Kim and J.S. Lim (Seoul, South Korea)	52-58
Vapo-liquid equilibria for the binary system of 2,3,3,3-tetrafluoroprop-1-ene (HFO-1234yf) + 1,1,1,2,3,3,3-heptafluoropropane (HFC-227ea) P. Hu, L.-X. Chen, W.-B. Zhu, L. Jia and Z.-S. Chen (Hefei, China)	59-61
A new biocompatible gentle aqueous biphasic system in cefalexin partitioning containing nonionic Tween 20 surfactant and three organic/inorganic different salts M. Taghavivand and G. Pazuki (Tehran, Iran)	62-71
Gas hydrate formation condition: Review on experimental and modeling approaches S. Shahnazar and N. Hasan (Kuala Lumpur, Malaysia)	72-85
Effect of 1-carboxymethyl-3-methylimidazolium chloride ionic liquid on thermodynamic and solubility of L-threonine in water at 298.15 K and atmospheric pressure M.T. Zafarani-Moattar and B. Asadzadeh (Tabriz, Iran)	86-95
Accurate prediction of the solubility parameter of pure compounds from their molecular structures T.A. Albahri (Safat, Kuwait)	96-103
Helmholtz function-based global phase stability test and its link to the isothermal-isochoric flash problem M. Castier (Doha, Qatar)	104-111
Effect of bivalent cation inorganic salts on isobaric vapor-liquid equilibrium of methyl acetate-methanol system J. Dhanalakshmi, P.S.T. Sai and A.R. Balakrishnan (Chennai, India)	112-119
Modeling of the thermodynamic properties of the mixtures: Prediction of the position of azeotropes for binary mixtures S. Fedali, H. Madani and C. Bougriou (Algeria)	120-127
Solubility and preferential solvation of sulfadiazine in methanol + water mixtures at several temperatures D.R. Delgado and F. Martínez (Bogotá D.C., Colombia)	128-138
Solid-liquid equilibrium prediction for binary mixtures of Ar, O ₂ , N ₂ , Kr, Xe, and CH ₄ using the LJ-SLV-EoS M. Campestrini, P. Stringari (Fontainebleau, France) and P. Arpentinier (Jouy-en-Josas, France)	139-147
Ionic liquids as entrainers for the separation of the ethanol/water system C. Tsanas, A. Tzani, A. Papadopoulos, A. Detsi and E. Voutsas	148-156
Water diffusivity in a mixed phase binary system: Effective water diffusion coefficients in tetradecanol-eicosanol mixtures D.R. Widmer, D.F. Calhoun, D. Thurston and A.M. Edwards (Granville, OH, USA)	157-166
Equation of state, elastic constants, and melting curve of solid neon using an effective two-body potential including quantum corrections M. Abbaspour and Z. Borzouie (Sabzevar, Iran)	167-174
Adsorption of He gas on the Ag _n nanoclusters: A molecular dynamic study H. Akbarzadeh and M. Mohammadzadeh (Sabzevar, Iran)	175-179
Thermodynamic solubility of pioglitazone HCl in polyethylene glycols 200, 400 or 600 + water mixtures at 303.2 and 308.2 K-Data report and modeling S. Soltanpour, F. Zohrabi and Z. Bastami (Zanjan, Iran)	180-184

(Contents continued on BMI)



(contents continued)

Phase equilibria of (water + butyric acid + butyl acetate) ternary systems at different temperatures K. Bayazıt, A. Göka, H. Uslu and Ş.İ. Kırbaşlar (Istanbul, Turkey).....	185–190
Critical temperatures and pressures of C ₄₀ , C ₄₄ , and C ₆₀ normal alkanes measured by the pulse-heating technique E.D. Nikitin and A.P. Popov (Ekaterinburg, Russia).....	191–195
One-parameter modified nonrandom two-liquid (NRTL) activity coefficient model S. Gebreyohannes, B.J. Neely (Stillwater, OK, USA) and K.A.M. Gasem (Stillwater, OK, USA and Laramie, WY, USA).....	196–205
Liquid–liquid equilibria of water + acetic acid + 2-ethyl hexyl acetate ternary system E. İnce and M. Lalikoglu (Istanbul, Turkey).....	206–211
Corrigendum	
Corrigendum to “Ternary (liquid–liquid) equilibrium data of furfuryl alcohol with organic solvents at $T = 298.2$ K: Experimental results and thermodynamic models” [Fluid Phase Equilib. 365 (2014) 88–96] T. Wongsawa (Bangkok, Thailand), M. Hronec, T. Soták (Bratislava, Slovakia), N. Leepipatpiboon, U. Pancharoen and S. Phatanasri (Bangkok, Thailand)	212–213