


F70



Volume 383, December 15 2014

ISSN 0378-3812

FLUID PHASE EQUILIBRIA

AN INTERNATIONAL JOURNAL

FLUID PHASE EQUILIBRIA

CONTENTS

(Abstracted/Index in: *Curr. Contents/Eng. Technol. Appl. Sci. Curr. Contents/Phys. Chem. Earth Sci., Sci. Cit. Index, Phys. Abstr., ASCA*, Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®)

Full Length Articles

Extraction of rhenium from sulfuric acid solution with used amine N235 D.-w. Fang, W.-j. Shan, Q. Yan, D. Li, L.-x. Xia (Shenyang, PR China) and S.-l. Zang (Shenyang, PR China and Fushun, PR China)	1-4
Experimental isobaric vapor-liquid equilibrium for ternary system of sec-butyl alcohol + sec-butyl acetate + <i>N,N</i> -dimethyl formamide at 101.3 kPa X. Zhang, Y. Liu, C. Jian, F. Wei and H. Liu (Tianjin, PR China)	5-10
Contribution to modeling the viscosity Arrhenius-type equation for some solvents by statistical correlations analysis R.B. Haj-Kacem (Nabeul, Tunisia), N. Ouerfelli (Tunis, Tunisia), J.V. Herráez (Burjassot, Spain), M. Guettari (Tunis, Tunisia), H. Hamda (La Marsa, Tunisia) and M. Dallel (Tunis, Tunisia)	11-20
The solid-liquid equilibrium, excess molar volume and refractive deviation properties of binary systems containing dimethyl carbonate, anisole and phenol S.-H. You, I.-Y. Jeong, I.-C. Hwang and S.-J. Park (Daejeon, Republic of Korea)	21-26
Solid-liquid equilibrium of ternary system sodium 1-naphthalenesulfonate + sodium 2-naphthalenesulfonate + water at (283.15, 303.15 and 323.15) K R. Li (Zhejiang, PR China and Hangzhou, PR China), G. Huang, H. Hu, C. Li (Zhejiang, PR China) and W. Jia (Zhejiang, PR China)	27-31
Apparent molar volumes and isentropic compressions of benzylalkylammonium ionic liquids in dimethylsulfoxide from 293.15 K to 328.15 K D. Keshapolla and R.L. Gardas (Chennai, India)	32-42
Liquid-liquid equilibria for ternary system water + toluene + benzaldehyde at (303.2-343.2) K H. Wang, Q. Wang (Hunan, PR China), Z. Xiong and C. Chen (Zhejiang, PR China)	43-48
Thermo-physical properties of ammonium-based ionic liquid + <i>N</i> -methyl-2-pyrrolidone mixtures at 298.15 K M. Usula (Monserrato, Italy), E. Matteoli (Pisa, Italy), F. Leonelli (Roma, Italy), F. Mocci (Monserrato, Italy and Stockholm, Sweden), F.C. Marincola (Monserrato, Italy), L. Gontran (Roma, Italy) and S. Porcedda (Monserrato, Italy)	49-54
Prediction of density, viscosity, and conductivity of the ternary aqueous solutions of piperidinium-based ionic liquids at different temperatures and atmospheric pressure using the data of their binary subsystems Y.-F. Chen, Y.-F. Hu, Z.-Y. Yang, J.-G. Qi, L.-Y. Yin, H.-R. Zhang, H.-Z. Huang and X.-M. Liu (Beijing, China)	55-71
Influence of the number, position and length of the alkyl-substituents on the solubility of water in pyridinium-based ionic liquids E.J. González and E.A. Macedo (Porto, Portugal)	72-77
Experimental study of the density and derived partial molar volumes of complex ternary water + 1-propanol + Li_2SO_4 mixtures at temperatures from (303 to 448) K and pressures up to 40 MPa I.M. Abdulagatov (Dagestan, Russia), L.A. Akhmedova-Azizova and N.D. Azizov (Baku, Azerbaijan)	78-93
Phase equilibrium data for the hydrogen sulphide + methane system at temperatures from 186 to 313 K and pressures up to about 14 MPa C. Coquelet, A. Valtz, P. Stringari, M. Popovic (Fontainebleau, France), D. Richon (Durban, South Africa and Aalto, Finland) and P. Mougin (Rueil-Malmaison, France)	94-99
The study of phase behavior of aqueous two-phase system containing $[\text{C}_n\text{mim}]^+ \text{BF}_4^-$ ($n = 2,3,4$) + $(\text{NH}_4)_2\text{SO}_4 + \text{H}_2\text{O}$ at different temperatures X. Tang, J. Han, Y. Hu, Y. Wang, Y. Lu, T. Chen and L. Ni (Zhenjiang, PR China)	100-107
Solubility modeling in three supercritical carbon dioxide, ethane and trifluoromethane fluids by one set of molecular descriptors R. Tabaraki and A. Toulabi (Ilam, Iran)	108-114
Fluctuational-thermodynamic interpretation of small angle X-ray scattering experiments in supercritical fluids V.B. Rogankov (Odessa, Ukraine)	115-125
The effect of the tetraalkylammonium salts on the protonation thermodynamics of the phytate anion C. Brett, R.M. Cigala, C. De Stefano, G. Lando and S. Sammartano (Messina, Italy)	126-133
COSMO-RS screening for efficient ionic liquid extraction solvents for NdCl_3 and DyCl_3 M. Grabda (Sendai, Japan and Zabrze, Poland), M. Panigrahi (Sendai, Japan), S. Oleszek (Sendai, Japan and Zabrze, Poland), D. Kozak (Sendai, Japan), F. Eckert (Leverkusen, Germany), E. Shibata and T. Nakamura (Sendai, Japan)	134-143

(Contents continued on BMI)



(contents continued)

Thermophysical properties of 1-butyl-3-methylimidazolium acetate over a wide range of temperatures and pressures J. Safarov (Rostock, Germany and Baku, Azerbaijan), M. Geppert-Rybczyńska (Katowice, Poland), I. Kul (Chester, PA, USA) and E. Hassel (Rostock, Germany).....	144–155
Generalized binary interaction parameters for the Peng–Robinson equation of state A.M. Abudour, S.A. Mohammad, R.L. Robinson Jr. (Stillwater, OK, USA) and K.A.M. Gasem (Laramie, OK, USA and WY, USA).....	156–173
Solubility of ammonium chloride in a $\text{MgCl}_2\text{-NH}_4\text{Cl-NH}_3\text{-H}_2\text{O}$ system at 298 K: Experiments, modeling and prediction H. Zhou, L. Huangfu, Y. Bao, X. Bai, R. Ma and Z. Sha (Tianjin, PR China).....	174–181
Separation of acetone and methanol azeotropic system using ionic liquid as entrainer W. Li, D. Sun, T. Zhang, S. Dai, F. Pan and Z. Zhang (Shenyang, China)	182–187

Letters to Editor

Comments on “isobaric (vapor + liquid) equilibria for three binary systems (toluene + anisole, <i>n</i> -butylbenzene + anisole, and guaiacol + anisole) at 101.33 kPa” J. Wisniak (Beer-Sheva, Israel), A. Mejía and H. Segura (Concepción, Chile).....	188–192
Reply to “Comments on ‘Isobaric (vapour + liquid) equilibria for three binary systems (toluene + anisole, <i>n</i> -butylbenzene + anisole, and guaiacol + anisole) at 101.33 kPa” H. Li, S. Xia, P. Ma, F.Y.Z. Yang and Y. Li (Tianjin, PR China)	193–196