


ПН  
F75/e

Volume 329, 1 October 2014

ISSN 0378-1127



ELSEVIER

# Forest Ecology and Management



**Special Section**  
**Restoring Working Forests in Human Dominated**  
**Landscapes of South Asia**

**Guest Editors**

Mark S. Ashton, Uromi M. Goodale, Kamaljit S. Bawa,  
J. David Neidel and Peter S. Ashton






ELSEVIER

Contents lists available at ScienceDirect

# Forest Ecology and Management

journal homepage: [www.elsevier.com/locate/foreco](http://www.elsevier.com/locate/foreco)

## Contents

### Special Section on Restoring Working Forests in Human Dominated Landscapes of South Asia

Guest Edited by Mark S. Ashton, Uromi M. Goodale, Kamaljit S. Bawa, J. David Neidel and Peter S. Ashton

|  |     |
|--|-----|
| Restoring working forests in human dominated landscapes of tropical south Asia<br>M.S. Ashton, U.M. Goodale, K.S. Bawa, J.D. Neidel and P.S. Ashton . . . . .  | 334 |
| Restoring working forests in human dominated landscapes of tropical South Asia: An introduction<br>M.S. Ashton (New Haven, USA), U.M. Goodale (Mengla, PR China), K.S. Bawa (Boston, USA), P.S. Ashton (Cambridge, USA) and<br>J. David Neidel (New Haven, USA) . . . . .  | 335 |
| Overcoming ecological barriers to tropical lower montane forest succession on anthropogenic grasslands: Synthesis and future<br>prospects<br>A.M.T.A. Gunaratne, C.V.S. Gunatilleke, I.A.U.N. Gunatilleke, H.M.S.P. Madawala (Peradeniya, Sri Lanka) and D.F.R.P. Burslem<br>(Aberdeen, UK) . . . . .  | 340 |
| Restoration of rain forest beneath pine plantations: A relay floristic model with special application to tropical South Asia<br>M.S. Ashton (New Haven, USA), C.V.S. Gunatilleke, I.A.U.N. Gunatilleke (Peradeniya, Sri Lanka), B.M.P. Singhakumara (Nugegoda,<br>Sri Lanka), S. Gamage (Peradeniya, Sri Lanka), T. Shibayama (Tokyo, Japan) and C. Tomimura (South Denpasar, Indonesia) . . . . . | 351 |
| Use and cultivation of plants that yield products other than timber from South Asian tropical forests, and their potential in forest<br>restoration<br>M.S. Ashton (United States), I.A.U.N. Gunatilleke, C.V.S. Gunatilleke (Sri Lanka), K.U. Tennakoon (Brunei Darussalam) and P.S. Ashton<br>(United States) . . . . .  | 360 |
| Altered stand structure and tree allometry reduce carbon storage in evergreen forest fragments in India's Western Ghats<br>A.M. Osuri (Bangalore, Mysore, India), V.S. Kumar (Bangalore, India) and M. Sankaran (Bangalore, India), (Leeds, UK) . . . . .  | 375 |
| The response of birds and mixed-species bird flocks to human-modified landscapes in Sri Lanka and southern India<br>E. Goodale (Mengla, China), S.W. Kotagama (Colombo, Sri Lanka), T.R.S. Raman, S. Sidhu (Mysore, India), U. Goodale (Mengla, China),<br>S. Parker (Cambridge, USA) and J. Chen (Mengla, China) . . . . .  | 384 |
| Cultural drivers of reforestation in tropical forest groves of the Western Ghats of India<br>S.A. Bhagwat (Oxford, United Kingdom), S. Nogué and K.J. Willis (Oxford, United Kingdom), (Bergen, Norway). . . . .   | 393 |
| <b>Regular Articles</b>  |     |
| How much carbon is sequestered during the restoration of tropical forests? Estimates from tree species in the Brazilian Atlantic forest<br>C.Y. Shimamoto (Curitiba, Brazil), P.C. Botosso (Colombo, Brazil) and M.C.M. Marques (Curitiba, Brazil) . . . . .   | 1   |
| Impact of silvicultural methods on vascular epiphytes (ferns, bromeliads and orchids) in a temperate forest in Oaxaca, Mexico<br>L.J. Bautista, A. Damon (Tapachula, Mexico), S. Ochoa-Gaona (Lerma Campeche, Mexico) and R.C. Tapia (Ixtlán de Juárez, Mexico) . . . . .  | 10  |
| Four-year response of underplanted American chestnut ( <i>Castanea dentata</i> ) and three competitors to midstory removal, root trenching,<br>and weeding treatments in an oak-hickory forest<br>E.D. Belair, M.R. Saunders (West Lafayette, United States) and B.G. Bailey (Sedro Woolley, United States). . . . .   | 21  |
| Long-term livestock grazing alters aspen age structure in the northwestern Great Basin<br>R.L. Beschta, J. Boone Kauffman (Corvallis, USA), D.S. Dobkin (Bend, USA) and L.M. Ellsworth (Corvallis, USA) . . . . .  | 30  |
| Sapling recruitment and mortality dynamics following partial harvesting in aspen-dominated mixedwoods in eastern Canada<br>A.K. Bose, B.D. Harvey and S. Brais (Rouyn-Noranda, Canada) . . . . .   | 37  |
| Drivers of lightning- and human-caused fire regimes in the Great Xing'an Mountains<br>T. Hu and G. Zhou (Beijing, China). . . . .  | 49  |
| Legacy effects of fire size and severity on forest regeneration, recruitment, and wildlife activity in aspen forests<br>H.Y. Wan, A.C. Olson, K.D. Muncey and S.B.St. Clair (Provo, USA). . . . .  | 59  |
| Cumulative effects of chronic deer browsing and clear-cutting on regeneration processes in second-growth white spruce stands<br>M. Barrette, L. Bélanger, L. De Grandpré and J.-C. Ruel (Québec, Canada) . . . . .   | 69  |
| Vertical stratification reduces competition for light in dense tropical forests<br>M. Laurans (Montpellier Cedex, France), B. Hérault (Kourou Cedex, French Guiana), G. Vieilledent and G. Vincent (Montpellier Cedex,<br>France). . . . .   | 79  |
| Oak savanna management strategies and their differential effects on vegetative structure, understory light, and flowering forbs<br>M.C. Lettow, L.A. Brudvig, C.A. Bahlai and D.A. Landis (East Lansing, USA) . . . . .  | 89  |

Contents continued on next page



|   |     |
|---|-----|
| Harvest residue removal and soil compaction impact forest productivity and recovery: Potential implications for bioenergy harvests<br>M.T. Curzon, A.W. D'Amato (Saint Paul, USA) and B.J. Palik (Grand Rapids, USA) . . . . .  | 99  |
| After long-term decline, are aspen recovering in northern Yellowstone?<br>L.E. Painter, R.L. Beschta (Corvallis, USA), E.J. Larsen (Stevens Point, USA) and W.J. Ripple (Corvallis, USA) . . . . .  | 108 |
| Vertical distribution and nitrogen content of epiphytic macrolichen functional groups in sub-boreal forests of central British Columbia<br>A. Kobylinski and A.L. Fredeen (Prince George, Canada) . . . . .   | 118 |
| Bromeliad rain: An opportunity for cloud forest management<br>T. Toledo-Aceves, J.G. García-Franco and F. López-Barrera (Xalapa, Mexico) . . . . .  | 129 |
| Response of tree regeneration to experimental gap creation and deer herbivory in north temperate forests<br>J.A. Forrester, C.G. Lorimer, J.H. Dyer, S.T. Gower and D.J. Mladenoff (Madison, USA) . . . . .   | 137 |
| Mediterranean <i>Quercus suber</i> wooded grasslands risk disappearance: New evidences from Sardinia (Italy)<br>I. Rossetti and S. Bagella (Sassari, Italy) . . . . .   | 148 |
| Climate-growth relationships for yellow-poplar across structural and site quality gradients in the southern Appalachian Mountains<br>T.L. Keyser (Asheville, United States) and P.M. Brown (Fort Collins, United States) . . . . .  | 158 |
| Effects of forest management on ground-dwelling beetles (Coleoptera; <i>Carabidae</i> , <i>Staphylinidae</i> ) in Central Europe are mainly mediated by changes in forest structure<br>M. Lange (Jena, Germany), M. Türke (Jena, Freising-Weihestephan, Germany), E. Pašalić (Jena, Germany), S. Boch (Bern, Switzerland), D. Hessenmöller (Jena, Germany), J. Müller (Potsdam, Germany), D. Prati, S.A. Socher, M. Fischer (Bern, Switzerland), W.W. Weisser and M.M. Gossner (Jena, Freising-Weihestephan, Germany) . . . . . | 166 |
| Assessment and prediction of above-ground biomass in selectively logged forest concessions using field measurements and remote sensing data: Case study in South East Cameroon<br>S. Gideon Neba, M. Kanninen (Finland), R. Eba'a Atyi and D.J. Sonwa (Yaounde, Cameroon) . . . . .   | 177 |
| The colonization of abandoned land by Spanish juniper: Linking biotic and abiotic factors at different spatial scales<br>B. Pías (Madrid, Spain), G. Escribano-Avila, E. Virgós (Móstoles, Spain), V. Sanz-Pérez (Madrid, Spain), A. Escudero (Móstoles, Spain) and F. Valladares (Móstoles, Madrid, Spain) . . . . .   | 186 |
| A new insight in the structure, composition and functioning of central African moist forests<br>A. Fayolle (Gembloux, Belgium), (Montpellier Cedex 5, France), N. Picard (Montpellier Cedex 5, France), J.-L. Doucet (Gembloux, Belgium), M. Swaine (Aberdeen, United Kingdom), N. Bayol (Mauguio, France), F. Bénédet and S. Gourlet-Fleury (Montpellier Cedex 5, France) . . . . .  | 195 |
| Assessing spatio-temporal rates, patterns and determinants of biological invasions in forest ecosystems. The case of <i>Acacia</i> species in NW Spain<br>L. Hernández, J. Martínez- Fernández (Madrid, Spain), I. Cañellas (Madrid, Palencia, Spain) and A.V. de la Cueva (Madrid, Spain) . . . . .  | 206 |
| Regional-scale application of lidar: Variation in forest canopy structure across the southeastern US<br>A.J. Hansen, L.B. Phillips (Bozeman, USA), R. Dubayah (College Park, USA), S. Goetz (Falmouth, USA) and M. Hofton (College Park, USA) . . . . .   | 214 |
| Combined effects of gap creation and deer exclusion on restoration of belowground systems of secondary woodlands: A field experiment in warm-temperate monsoon Asia<br>M. Suzuki and E. Ito (Japan) . . . . .   | 227 |
| A critical review of forest biomass estimation models, common mistakes and corrective measures<br>G.W. Sileshi (Lilongwe, Malawi) . . . . .   | 237 |
| Variations in the live biomass and carbon pools of <i>Abies georgei</i> along an elevation gradient on the Tibetan Plateau, China<br>G. Wang, F. Ran, R. Chang, Y. Yang, J. Luo and F. Jianrong (Chengdu, PR China) . . . . .   | 255 |
| Fire severity, size, and climate associations diverge from historical precedent along an ecological gradient in the Pinaleno Mountains, Arizona, USA<br>C.D. O'Connor, D.A. Falk, A.M. Lynch and T.W. Swetnam (Tucson, United States) . . . . .   | 264 |
| Repeated harvest residue removal reduces <i>E. globulus</i> productivity in the 3rd rotation in south-western Australia<br>D.S. Mendham (Sandy Bay, Australia), G.N. Ogden, T. Short, T.M. O'Connell, T.S. Grove and S.J. Rance (Floreat, Australia) . . . . .  | 279 |
| Quantity and quality of dissolved organic carbon released from coarse woody debris of different tree species in the early phase of decomposition<br>A. Bantle, W. Borken (Bayreuth, Germany), R.H. Ellerbrock (Müncheberg, Germany), E.D. Schulze (Jena, Germany), W.W. Weisser (Freising-Weihestephan, Germany) and E. Matzner (Bayreuth, Germany) . . . . .   | 287 |
| Impact of the first recorded outbreak of the Douglas-fir tussock moth, <i>Orgyia pseudotsugata</i> , in southern California and the extent of its distribution in the Pacific Southwest region<br>T.W. Coleman (San Bernardino, United States), M.I. Jones (Davis, United States), B. Courtial (Orléans, France), A.D. Graves (Albuquerque, United States), M. Woods (Davis, United States), A. Roques (Orléans, France) and S.J. Seybold (Davis, United States) . . . . .  | 295 |
| A compatible system of biomass equations for three conifer species in Northeast, China<br>L. Dong (Harbin, PR China), L. Zhang (Syracuse, USA) and F. Li (Harbin, PR China) . . . . .   | 306 |
| Understory species regulate litter decomposition and accumulation of C and N in forest soils: A long-term dual-isotope experiment<br>Y. Qiao (Harbin, China), (Davis, USA), S. Miao (Harbin, China), L.C.R. Silva and W.R. Horwath (Davis, USA) . . . . .   | 318 |
| Is facilitation a promising strategy for cloud forest restoration?<br>M.d.I.L. Avendaño-Yáñez, L.R. Sánchez-Velásquez (Xalapa, Mexico), J.A. Meave (México, Mexico) and M.d.R. Pineda-López (Xalapa, Mexico) . . . . .  | 328 |